

Literatur

Literatur zu Kapitel 2 Trainingsformen/Komponenten der Bewegungs- und Sporttherapie

1. Pedersen BK, Saltin B. Exercise as medicine – evidence for prescribing exercise as therapy in 26 different chronic diseases. *Scand J Med Sci Sports* 2015; 25, Suppl 3: 1–72.
2. DVGS online: <http://www.dvgs.de/dvgs-verband/sport-bewegungstherapie/definition.html>; veröffentlicht am 10. April 2014.
3. Hollmann W, Hettinger T. Sportmedizin. Grundlagen für Arbeit, Trainings- und Präventivmedizin. Stuttgart: Schattauer, 2009.
4. Diemer F, Sutor V. Praxis der medizinischen Trainingstherapie I: Lendenwirbelsäule, Sakroiliakgelenk und untere Extremität. Stuttgart: Thieme, 2011.
5. Weineck J. Sportbiologie (10., überarbeitete und erweiterte Auflage). Balingen: Spitta, 2010.
6. Kindermann W. Anaerobe Schwelle. *Dtsch Z Sportmed* 2004; 55: 161–62.
7. Paffenbarger RS Jr, Wing AL, Hyde RT. Physical activity as an index of heart attack risk in college alumni. *Am J Epidemiol* 1978; 142: 889–903.
8. Haskell WL, Lee IM, Pate RR, Powell KE, Blair SN, Franklin BA et al. Physical activity and public health: updated recommendation for adults from the American College of Sports Medicine and the American Heart Association. *Med Sci Sports Exerc* 2007; 39: 1423–34.
9. Moyna NM, Thompson PD. The effect of physical activity on endothelial function in man. *Acta Physiol Scand* 2004; 180: 113–23.
10. Daugaard JR, Richter EA. Relationship between muscle fibre composition, glucose transporter protein 4 and exercise training: possible consequences in non-insulin-dependent diabetes mellitus. *Acta Physiol Scand* 2001; 171: 267–76.
11. Prabhakaran B, Dowling EA, Branch JD, Swain DP, Leutholtz BC. Effect of 14 weeks of resistance training on lipid profile and body fat percentage in premenopausal women. *Br J Sports Med* 1999; 33: 190–5.
12. Predel HG. Bluthochdruck und Sport. *Dtsch Z Sportmed* 2007; 58: 328–33.
13. Halloway S, Wilbur J, Schoeny ME, Arfanakis K. Effects of Endurance-Focused Physical Activity Interventions on Brain Health: A Systematic Review. *Biol Res Nurs* 2016; Jul 29. pii: 1099800416660758.
14. Garber CE, Blissmer B, Deschenes MR, Franklin BA, Lamonte MJ et al. American College of Sports Medicine position stand. Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults: guidance for prescribing exercise. *Med Sci Sports Exerc* 2011; 43: 1334–59.
15. Wahl P, Hägele M, Zinner C, Bloch W, Mester J. High Intensity (HIT) für die Verbesserung der Ausdauerleistungsfähigkeit von Normalpersonen und im Präventions- & Rehabilitationsbereich. *Wien Med Wochenschr* 2010; 160: 627–36.
16. Vogiatzis I, Terzis G, Nanas S, Stratakis G, Simoes DC, Georgiadou O et al. Skeletal muscle adaptations to interval training in patients with advanced COPD. *Chest* 2005; 128: 3838–45.
17. Gartner-Schmidt JL, Shembel AC, Zullo TG, Rosen CA. Development and validation of the Dyspnea Index (DI): a severity index for upper airway-related dyspnea. *J Voice: official journal of the Voice Foundation* 2014; 28: 775–82.
18. Donnelly JE, Blair SN, Jakicic JM, Manore MM, Rankin JW, Smith BK; American College of Sports Medicine. American College of Sports Medicine Position Stand. Appropriate physical activity intervention strategies for weight loss and prevention of weight regain for adults. *Med Sci Sports Exerc* 2009; 41: 459–71.
19. Braith RW, Stewart KJ. Resistance exercise training: its role in the prevention of cardiovascular disease. *Circulation* 2006; 113: 2642–50.
20. Kelley G. Dynamic resistance exercise and resting blood pressure in adults: a meta-analysis. *J Appl Physiol* 1997; 82: 1559–65.
21. Fleck S, Dean L. Resistance-training experience and the pressor response during resistance exercise. *J Appl Physiol* 1997; 63: 116–20.
22. Sherrington C, Tiedemann A, Fairhall N, Close JC, Lord SR. Exercise to prevent falls in older adults: an updated meta-analysis and best practice recommendations. *N S W Public Health Bull* 2011; 22: 78–83.
23. Decoster LC, Cleland J, Altieri C, Russell P. The effects of hamstring stretching on range of motion: a systematic literature review. *J Orthop Sports Phys Ther* 2005; 35: 377–87.
24. Costa PB, Graves BS, Whitehurst M, Jacobs PL. The acute effects of different durations of static stretching on dynamic balance performance. *J Strength Cond Res* 2009; 23: 141–7.
25. Klee A. Methoden und Wirkungen des Dehnungstrainings. Die Ruhespannungs-Dehnungskurve – ihre Erhebung beim M. rectus femoris und ihre Veränderung im Rahmen kurzfristiger Treatments. Habilitationsschrift. Schorndorf: Verlag K. Hofmann, 2003.

26. Feland JB, Myrer JW, Schulthies SS, Fellingham GW, Measom GW. The effect of duration of stretching of the hamstring muscle group for increasing range of motion in people aged 65 years or older. *Phys Ther* 2001; 81: 1110–7.
27. Stehle P. BISP-Expertise Sensomotorisches Training – Propriozeptives Training. Köln: Sportverlag Strauß, 2009.
28. Zech A, Hübscher M, Vogt L, Banzer W, Hänsel F, Pfeifer K. Balance training for neuromuscular control and performance enhancement: a systematic review. *J Athl Train* 2010; 45: 392–403.
29. Hirtz P. Koordinative Fähigkeiten – Kennzeichnung, Altersgang und Beeinflussungsmöglichkeiten. *Medizin und Sport* 1981; 21: 348–51.
30. Kolb M. Spiele für den Herz- und Alterssport. Aachen: Meyer & Meyer Sport, 2007.
31. Schüle K, Huber G. Grundlagen der Sport- und Bewegungstherapie: Prävention, ambulante und stationäre Rehabilitation. Köln: Deutscher Ärzte-Verlag, 2012.
32. Fuchs R. Das MoVo-Modell als theoretische Grundlage für Programme der Gesundheitsverhaltensänderung. In: R. Fuchs, W. Göhner & H. Seelig (Hrsg.) *Aufbau eines körperlich aktiven Lebensstils*. Göttingen: Hogrefe, 2007: S. 317–25.
33. Geidl W, Hofmann J, Göhner W, Sudeck G, Pfeifer K. Verhaltensbezogene Bewegungstherapie – Bindung an einen körperlich aktiven Lebensstil. *Behaviour-Orientated Exercise Therapy – Initiating and Maintaining a Physically Active Lifestyle*. Rehabilitation 2012; 51: 259–68.
34. Mühlig S. Allgemeine Trainerqualifikationen zur Patientenschulung: Defizite und Qualitätskriterien. *Praxis klinische Verhaltensmedizin und Rehabilitation* 2007; 20: 74–9.
35. Göhner W, Fuchs R. Änderung des Gesundheitsverhaltens: MoVo-Gruppenprogramme für körperliche Aktivität und gesunde Ernährung. Göttingen: Hogrefe, 2007.

Literatur zu Kapitel 3 Adaptation des Körpers an Bewegung

1. Calabrese EJ, Baldwin LA. Defining hormesis. *Hum Exp Toxicol* 2002; 21: 91–7.
2. Hellsten Y, Nyberg M. Cardiovascular Adaptations to Exercise Training. *Compr Physiol* 2015; 6: 1–32.
3. Kindermann W. Physiologische Anpassungen des Herz-Kreislauf-Systems an körperliche Belastung. In: Kindermann W, Dickhuth HH, Nieß A, Röcker K, Urhausen A (Hrsg): *Sportkardiologie – körperliche Aktivität bei Herzerkrankungen*. Steinkopff, Darmstadt, 2. Auflage, 2007, 1–20.
4. Flück M, Hoppeler H. Molecular basis of skeletal muscle plasticity – from gene to form and function. *Rev Physiol Biochem Pharmacol*. 2003; 146:159–216. Epub 2003 Jan 14. Review.
5. Folland JP, Williams AG. The adaptations to strength training : morphological and neurological contributions to increased strength. *Sports Med* 2007; 37: 145–68. Review.
6. Booth FW, Thomason DB. Molecular and cellular adaptation of muscle in response to exercise: perspectives of various models. *Physiol Rev* 1991; 71: 541–85.
7. Mooren FC, Krüger K. Apoptotic lymphocytes induce progenitor cell mobilization after exercise. *J Appl Physiol* (1985) 2015; 119: 135–9.
8. Kjaer M, Magnusson P, Krogsgaard M, Boysen Møller J, Olesen J, Heinemeier K et al. Extracellular matrix adaptation of tendon and skeletal muscle to exercise. *J Anat* 2006; 208: 445–50.
9. Eckstein F, Hudelmaier M, Putz R. The effects of exercise on human articular cartilage. *J Anat* 2006; 208: 491–512.
10. Hannukainen JC, Borra R, Linderborg K, Kallio H, Kiss J, Lepomäki V et al. Liver and pancreatic fat content and metabolism in healthy monozygotic twins with discordant physical activity. *J Hepatol* 2011; 54: 545–552.
11. Cerdá B, Pérez M, Pérez-Santiago JD, Tornero-Aguilera JF, González-Soltero R, Larrosa M. Gut Microbiota Modification: Another Piece in the Puzzle of the Benefits of Physical Exercise in Health? *Front Physiol* 2016; 7: 51.
12. Stanford KI, Roeland J. W. Middelbeek. Goodyear LJ. Exercise Effects on White Adipose Tissue: Beiging and Metabolic Adaptations. *Diabetes*; 64: 2361–2368.
13. Ruscheweyh R, Willemer C, Krüger K, Duning T, Warnecke T, Sommer J et al. Physical activity and memory functions: an interventional study. *Neurobiol Aging* 2011; 32: 1304–19.
14. Eyre H, Baune BT. Neuroimmunological effects of physical exercise in depression. *Brain Behav Immun* 2012; 26: 251–66.
15. Bayod S, Del Valle J, Canudas AM, Lalanza JF, Sanchez-Roige S, Camins A, Escorihuela RM, Pallàs M. Long-term treadmill exercise induces neuroprotective molecular changes in rat brain. *J Appl Physiol* (1985) 2011; 111: 1380–90.
16. Wrann CD, White JP, Salogiannis J, Laznik-Bogoslavski D, Wu J, Ma D et al. Exercise induces hippocampal BDNF through a PGC-1 α /FNDC5 pathway. *Cell Metab* 2013; 18: 649–59.
17. Radák Z, Kaneko T, Tahara S, Nakamoto H, Ohno H, Sasvári M, Nyakas C, Goto S. The effect of exercise training on oxidative damage of lipids, proteins, and DNA in rat skeletal muscle: evidence for beneficial outcomes. *Free Radic Biol Med* 1999; 27: 69–74.
18. Powers SK, Radak Z, Ji LL. Exercise-induced oxidative stress: past, present and future. *J Physiol* 2016 Feb 19.

19. Fehrenbach E, Northoff H. Free radicals, exercise, apoptosis, and heat shock proteins. *Exerc Immunol Rev* 2001; 7: 66–89. Review.
20. McGee SL, Hargreaves M. Histone modifications and exercise adaptations. *J Appl Physiol* (1985) 2011; 110: 258–63.
21. Luttrop K, Nordfors L, Ekström TJ, Lind L. Physical activity is associated with decreased global DNA methylation in Swedish older individuals. *Scand J Clin Lab Invest* 2013; 73: 184–5.
22. Meurer S, Krüger K, Mooren FC. MicroRNAs and Exercise. *Dtsch Z Sportmed* 2016; 67: 27–34.

Literatur zu Kapitel 4 Methoden der Trainingssteuerung

1. Holland AE et al. An official European Respiratory Society/American Thoracic Society technical standard: field walking tests in chronic respiratory disease. *Eur Respir J* 2014; 44: 1428–46.
2. Enright PL. The six-minute walk test. *Respir Care* 2003; 48: 783–5.
3. American College of Sports. ACSM's resource manual for guidelines for exercise testing and prescription. 5th ed ed. 2006, Philadelphia: Lippincott Williams und Wilkins.
4. Borg G. Physical performance an perceived exertion. *Studia Psychologia et Paedagogica Invest XI* 1962.
5. Borg G. Anstrengungsempfinden und körperliche Arbeit. *Dtsch Arztebl* 2004; 15: 101.
6. Abbiss C, Laursen P. Models to explain fatigue during prolonged endurance cycling. *Sports Med* 2005; 35: 865–98.
7. Noble BJ, Robertson RJ. Perceived exertion. *Human Kinetics*. XV, Champaign 1996: 320.
8. Feigenbaum MS. Resistance training for health and rehabilitation, in *Human Kinetics II*, Champaign. 2001.
9. Graves JG, Franklin BA. Resistance training for health and rehabilitation. *Human Kinetics II*. Champaign 2001.
10. Löllgen H, Erdmann E. Ergometrie. Berlin, Heidelberg: Springer-Verlag, 2000.
11. Samitz G, Mensink G. Körperliche Aktivität in Prävention und Therapie – Evidenzbasierter Leitfaden für Klinik und Praxis. München: Hans Marseille Verlag, 2002.

Literatur zu Kapitel 5 Sportärztliche Vorsorgeuntersuchung

1. Löllgen H, Löllgen D. Risikoreduzierung durch körperliche Aktivität. *Der Internist* 2012; 53: 20–9.
2. Löllgen H, Böckenhoff A, Knapp G. Primary prevention by physical activity: an updated meta-analysis with different intensity categories. *Int. J Sports Med*. 2009; 30: 213–24.
3. Samitz G, Egger M, Zwahlen M. Domains of physical activity and all-cause mortality: systematic review and dose response meta-analysis of cohort studies. *Int J Epidemiol* 2011; 40: 1382–400.
4. Priori SG, Zipes DP. Sudden cardiac death. Malden, Massachusetts: Blackwell 2006.5.
5. Löllgen H, Gerke R, Steinberg T. Der kardiale Zwischenfall im Sport. *Dtsch Arztebl* 2006; 103: A1617–21.6.
6. Maron BJ, Doerer JJ, Haas TS, Tierney DM, Mueller FO. Sudden death in young competitive athletes. *Circulation* 2009; 119: 1085–92.
7. Tester DJ, Ackerman MJ. The role of molecular autopsy in unexplained sudden cardiac death. *Curr Opin Cardiol* 2006; 21:166–72.
8. Drezner JA. Contemporary approaches to the identification of athletes at risk for sudden cardiac death. *Curr Opin Cardiol* 2008; 494–501.
9. Klot S, Mittleman MA, Dockery DW, Heier M, Meisinger C et al. Intensity of physical exertion and triggering of myocardial infarction: a case-crossover study *Eur Heart J* 2008; 29: 1881–8.
10. Marti B. Plötzliche Todesfälle an Schweizer Volksläufen 1978–1987. *Schweiz Med Wschr* 1989; 119: 473–82.
11. Harmon KG, Asif IM, Klossner D, Drezner JA. Incidence of sudden cardiac death in national collegiate athletic association athletes. *Circulation* 2011; 123: 1594–600.
12. Corrado D, Basso C, Rizzoli G, Schiavon M, Thiene G. Does sports activity enhance the risk of sudden death in adolescents and young adults? *J Am Coll Cardiol* 2003; 42: 959–63.
13. Drory Y, Turetz Y, Hiss Y, Lev B, Fisman EZ et al. Sudden unexpected death in persons less than 40 years of age. *Am J Cardiol* 1991; 68: 1388–92.
14. Maron BJ. How should we screen competitive athletes for cardiovascular disease? *Europ Heart J* 2005; 8: 516–24.
15. Papadakis M, Whyte G, Sharma S. Preparticipation screening for cardiovascular abnormalities in young competitive athletes. *Brit Med J* 2008; 337: 806–11.
16. Sen-Chowdhry S, Syrris P, Prasad SK, Hughes SE, Merrifield R et al. Left-dominant arrhythmogenic cardiomyopathy: an under-recognized clinical entity. *J Am Coll Cardiol* 2008; 52: 2175–87.
17. Dickhuth HH, Mayer F, Röcker K, Berg A (eds.) *Sportmedizin für Ärzte*. Köln: Deutscher Ärzte-Verlag 2007.
18. Löllgen H: Kardiopulmonale Funktionsdiagnostik, 4th ed., Nürnberg: Novartis 2005.

19. Myerburg RJ, Vetter VL. Electrocardiograms should be included in preparticipation screening of athletes. *Circulation* 2007; 116: 2616–26.
20. Fuller CM, McNulty CM, Spring DA, Arger KM, Bruce SS et al. Prospective screening of 5615 high school athletes for risk of sudden cardiac death. *Med Sci Sports Exerc* 1997; 29: 1131–8.
21. Pelliccia A, Maron BJ: Preparticipation cardiovascular evaluation of the competitive athlete: perspectives from the 30-year Italian experience. *Am J Cardiol* 1995; 75: 827–9.
22. Maron BJ, Douglas PS, Graham TP, Nishimura RA, Thompson PD. Task Force 1: Preparticipation screening and diagnosis of cardiovascular disease in athletes. *J Am Coll Cardiol*. 2005; 45: 1322–6.
23. Maron BJ, Araújo CG, Thompson PD, Fletcher GF, de Luna AB et al. Recommendations for preparticipation screening and the assessment of cardiovascular disease in masters athletes: an advisory for healthcare professionals from the working groups of the World Heart Federation, the International Federation of Sports Medicine, and the American Heart Association Committee on Exercise, Cardiac Rehabilitation, and Prevention, *Circulation* 2001; 103: 327–34.
24. Corrado D, Basso C, Schiavon M, Pelliccia A, Thiene G. Pre-Participation screening of young competitive athletes for prevention of sudden cardiac death. *J Am Coll Cardiol* 2008; 52: 1981–9.
25. Pelliccia A, Zipes DP, Maron BJ. Bethesda conference #36 and the European Society of Cardiology consensus recommendations revisited: A comparison of U.S. and European criteria for eligibility and disqualification of competitive athletes with cardiovascular abnormalities *J Am Coll Cardiol* 2008; 52: 1990–6.
26. Bernhardt, DT, Roberst WO (eds.) : Preparticipation physical evaluation. 4th ed. Am Acad Pediatrics, 2010.
27. Hoffmann A, Isler R. Kardiale Beurteilung der Sporttauglichkeit. *Schweiz Med Forum* 2007; 7: 889–94.
28. Löllgen H, Leyk D, Hansel J. The Pre-Participation Examination for Leisure Time Physical Activity. General medical and Cardiological Issues. Review. *Dtsch Arztebl Int* 2010; 107: 1–8, I-II.
29. Löllgen H, Björjesson M, Cumiskey J, Bachl N, Debruyne A: The Pre-Participation Examination in Sports: EFSMA Statement on ECG for Pre-Participation Examination. *Dtsch Zschr Sportmed* 2015; 66: 151–155.
30. DGSP: Leitlinie Vorsorgeuntersuchung im Sport. Frankfurt 2007.
31. Wilson MG, Basavarajiah S, Whyte GP, Cox S, Loosemore M, Sharma S. Efficacy of personal symptom and family history questionnaire when screening for inherited cardiac pathologies: the role of electrocardiography. *Br J Sports Med* 2008; 42: 207–11.
32. Steinvil A, Chundadze T, Zeltser D, Rogowski O, Halkin A et al. Mandatory electrocardiographic screening of athletes to reduce their risk for sudden death proven fact or wishful thinking? *J Am Coll Cardiol* 2011; 57: 1291–6.
33. Pelliccia A, Saner H. Participation in leisure-time physical activities and competitive sports in patients with cardiovascular diseases: How to get the benefits without risks. *Europ J Cardiovasc Prev Rehab* 2005; 12: 315–7.
34. Trappe H-J, Löllgen H. Leitlinien zur Ergometrie. *Z Kardiol* 2000; 89: 821–37.
35. Gibbons RJ, Balady GJ, Bricker JT: ACC/AHA 2002 guidelines update for exercise testing. Summary article. *Circulation* 2002; 883–92.
36. Corrado D, Pelliccia A, Bjørnstad HH, Vanhees L, Biffi A et al. Cardiovascular pre-participation screening of young competitive athletes for prevention of sudden death: proposal for a common European protocol. Consensus Statement of the Study Group of Sport Cardiology of the Working Group of Cardiac Rehabilitation and Exercise Physiology and the Working Group of Myocardial and Pericardial Diseases of the European Society of Cardiology. *Eur Heart J* 2005; 26: 516–24.
37. ACSM guidelines for exercise testing and prescription, 2014, 9th ed. Philadelphia: Lippincott (American College of Sports Medicine (ACSM)).
38. Löllgen H, Erdmann E, Gitt A. Ergometrie. 3rd ed. Heidelberg: Springer 2009. ISBN-10: 3540927298.
39. Bachl N., Schwarz W, Zeibig J. Fit ins Alter: Mit richtiger Bewegung jung bleiben, Springer 2006, ISBN-10: 3211235232, S 72–80.
40. Maron BJ, Zipes DP: 36th Bethesda Conference: Eligibility recommendations for competitive athletes with cardiovascular abnormalities. *J Am Coll Cardiol* 2005; 45: 1312–75.
41. Warburton DER, Jamnik VK, Bredin SSD, McKenzie DC, Stone J et al. Evidence-based risk assessment and recommendations for physical activity clearance: an introduction. *Appl Physiol Nutr Metab* 2011; 36: 1–2.

Literatur zu Kapitel 6 Sport und Medikation

1. Mooren F C, Stein B. Schadet Marathonlaufen dem Gastrointestinaltrakt? *Dtsch Z Sportmed* 2011; 62, 9: 304–309.
2. Kargotich S, Goodman C, Keast D, Morton AR. The influence of exercise-induced plasma volume changes on the interpretation of biochemical parameters used for monitoring exercise, training and sport. *Sports Med* 1998; 26: 101–17.
3. Ardies CM, Morris GS, Erickson CK, Farrar RP. Both acute and chronic exercise enhance in vivo ethanol clearance in rats. *J Appl Physiol* (1985) 1989; 66: 555–60.

4. Predel HG. Bluthochdruck und Sport. *Dt Zeitschr Sportmedizin* 2007; 58: 328–333.
5. Ciccoira M, Zanolla L, Rossi A, Golia G, Franceschini L, Brighetti G, Marino P, Zardini P. Long-term, dose-dependent effects of spironolactone on left ventricular function and exercise tolerance in patients with chronic heart failure. *J Am Coll Cardiol* 2002 40: 304–10.
6. Onder G, Vedova CD, Pahor M. Effects of ACE inhibitors on skeletal muscle. *Curr Pharm Des* 2006; 12: 2057–64.
7. Onder G, Penninx BW, Balkrishnan R, Fried LP, Chaves PH, Williamson J, Carter C, Di Bari M, Guralnik JM, Pahor M. Relation between use of angiotensin-converting enzyme inhibitors and muscle strength and physical function in older women: an observational study. *Lancet* 2002; 359: 926–30.
8. Wonisch M, Hofmann P, Fruhwald FM, Kraxner W, Hödl R, Pokan R, Klein W. Influence of beta-blocker use on percentage of target heart rate exercise prescription. *Eur J Cardiovasc Prev Rehabil*. 2003; 10: 296–301.
9. Steinshamn S, Sandsund M, Sue-Chu M, Bjørmer L. Effects of montelukast and salmeterol on physical performance and exercise economy in adult asthmatics with exercise-induced bronchoconstriction. *Chest* 2004; 126: 1154–60.
10. Glass DJ. Signaling pathways perturbing muscle mass. *Curr Opin Clin Nutr Metab Care* 2010; 13: 225–9.
11. Mooren FC. Diabetes mellitus Typ 2 und Sport – Blutzuckerkontrolle durch Ausdauer- und Krafttraining. *Kliniker* 2013; 42: 406–410.
12. Burkhard PR. Acute and subacute drug-induced movement disorders. *Parkinsonism Relat Disord* 2014; 20 Suppl 1: S108–12.
13. Rojas-Fernandez CH, Seymour N, Brown SG. Helping pharmacists to reduce fall risk in long-term care: A clinical tool to facilitate the medication review process. *Can Pharm J (Ott)* 2014; 147: 171–8.

Literatur zu Kapitel 7 Mortalität und Lebenserwartung

1. LaFortune L, Martin S, Kelly S, Kuhn I, Remes O et al. Behavioural risk factors in mid-life associated with successful ageing, disability, dementia and frailty in later life: a rapid systematic review. *PLoS One* 2016; 11: e0144405.
2. Rezende LF, Sá TH, Mielke GI, Viscondi JY, Rey-López JP, Garcia LM. All-cause mortality attributable to sitting time: Analysis of 54 countries worldwide. *Am J Prev Med* 2016; 51: 253–63.
3. Hupin D, Roche F, Gremeaux V, Chatard JC, Oriol M et al. Even a low-dose of moderate-to-vigorous physical activity reduces mortality by 22 % in adults aged ≥60 years: a systematic review and meta-analysis. *Br J Sports Med* 2015; 49: 1262–7.
4. Stewart RA, Benatar J, Maddison R. Living longer by sitting less and moving more. *Curr Opin Cardiol* 2015; 30: 551–7.
5. Stamatakis E, Rogers K, Ding D, Berrigan D, Chau J et al. All-cause mortality effects of replacing sedentary time with physical activity and sleeping using an isotemporal substitution model: a prospective study of 201,129 mid-aged and older adults. *Int J Behav Nutr Phys Act* 2015; 12: 121.
6. Schmid D, Ricci C, Baumeister SE, Leitzmann MF. Replacing sedentary time with physical activity in relation to mortality. *Med Sci Sports Exerc* 2016; 48: 1312–9.
7. Ekelund U, Steene-Johannessen J, Brown WJ, Fagerland MW, Owen N et al. Does physical activity attenuate, or even eliminate, the detrimental association of sitting time with mortality? A harmonised meta-analysis of data from more than 1 million men and women. *Lancet* 2016; 388: 1302–10.
8. Reimers CD, Reimers AK, Knapp G. Prävention durch körperliche Aktivität und Sport. In: Mewes N, Reimers CD, Knapp G (Hrsg.) *Prävention und Therapie durch Sport. Band 1: Grundlagen*. Elsevier Urban & Fischer, München 2015, S. 3–87.
9. Rijk JM, Roos PR, Deckx L, van den Akker M, Buntinx F. Prognostic value of handgrip strength in people aged 60 years and older: A systematic review and meta-analysis. *Geriatr Gerontol Int* 2016; 16: 5–20.
10. Strand BH, Cooper R, Bergland A, Jørgensen L, Schirmer H et al. The association of grip strength from midlife onwards with all-cause and cause-specific mortality over 17 years of follow-up in the Tromsø Study. *J Epidemiol Community Health*. 2016 May 26. pii: jech-2015–206776. doi: 10.1136/jech-2015–206776. [Epub ahead of print]
11. Moore SC, Patel AV, Matthews CE, Berrington de Gonzalez A, Park Y et al. Leisure time physical activity of moderate to vigorous intensity and mortality: a large pooled cohort analysis. *PLoS Med* 2012; 9: e1001335.
12. Loprinzi PD. Accumulated short bouts of physical activity are associated with reduced premature all-cause mortality: implications for physician promotion of physical activity and revision of current US government physical activity guidelines. *Mayo Clin Proc* 2015; 90: 1164–9.
13. Ferrucci L, Izmirlian G, Leveille S, Phillips CL, Corti M-C et al. Smoking, physical activity, and active life expectancy. *Am J Epidemiol* 1999; 149: 645–53.
14. Fraser GE, Shavlik DJ. Ten years of life. Is it a matter of choice? *Arch Intern Med* 2001; 161: 1645–52.
15. Janssen I, Carson V, Lee I-M, Katzmarzyk PT, Blair SN. Years of life gained due to leisure-time physical activity in the United States. *Am J Prev Med*. Author manuscript; available in PMC 2014 January 1.

16. Jonker JT, De Laet C, Franco OH, Peeters A, Mackenbach J, Nusselder WJ. Physical activity and life expectancy with and without diabetes. *Diabetes Care* 2006; 29: 38–43.
17. Nusselder WJ, Franco OH, Peeters A, Mackenbach JP. Living healthier for longer: Comparative effects of three heart-healthy behaviors on life expectancy with and without cardiovascular disease. *BMC Public Health* 2009; 9: 487.
18. O'Doherty MG, Cairns K, O'Neill V, Lamrock F, Jørgensen T et al. Effect of major lifestyle risk factors, independent and jointly, on life expectancy with and without cardiovascular disease: results from the Consortium on Health and Ageing Network of Cohorts in Europe and the United States (CHANCES). *Eur J Epidemiol* 2016; 31: 455–68.
19. Paffenbarger Jr. RS, Hyde RT, Hsieh C-C, Wing AL. Physical activity, other life-style patterns, cardiovascular disease and longevity. *Acta Med Scand Suppl* 1986; 711: 85–91.
20. Paffenbarger Jr. RS, Kampert JB, Lee IM, Hyde RT, Leung RW, Wing AL. Changes in physical activity and other lifeway patterns influencing longevity. *Med Sci Sports Exerc* 1994; 26: 857–65.
21. Pekkanen J, Marti B, Nissinen A, Tuomilehto J, Punsar S, Karvonen MJ. Reduction of premature mortality by high physical activity: a 20-year follow-up of middle-aged Finnish men. *Lancet* 1987; 1: 1473–7.
22. Wen CP, Wai JP, Tsai MK, Yang YC, Cheng TY et al. Minimum amount of physical activity for reduced mortality and extended life expectancy: a prospective cohort study. *Lancet* 2011; 378: 1244–53.
23. Lemez S, Baker J. Do elite athletes live longer? A systematic review of mortality and longevity in elite athletes. *Sports Med Open* 2015; 1: 16.
24. Kettunen JA, Kujala UM, Kaprio J, Bäckmand H, Peltonen M et al. All-cause and disease-specific mortality among male, former elite athletes: an average 50-year follow-up. *Br J Sports Med* 2015; 49: 893–7.
25. Garatachea N, Santos-Lozano A, Sanchis-Gomar F, Fiuza-Luces C, Pareja-Galeano H et al. Elite athletes live longer than the general population: a meta-analysis. *Mayo Clin Proc* 2014; 89: 1195–200.
26. Kim JH, Baggish AL. Physical activity, endurance exercise, and excess – Can one overdose? *Curr Treat Options Cardiovasc Med.* 2016; 18: 68.
27. Karvinen S, Waller K, Silvennoinen M, Koch LG, Britton SL et al. Physical activity in adulthood: genes and mortality. *Sci Rep* 2015; 5: 18259.
28. Garatachea N, Pareja-Galeano H, Sanchis-Gomar F, Santos-Lozano A, Fiuza-Luces C et al. Exercise attenuates the major hallmarks of aging. *Rejuvenation Res* 2015; 18: 57–89.

Literatur zu Kapitel 8 Prähabilitation

1. Santa Mina D, Scheede-Bergdahl C, Gillis C, Carli F. Optimization of surgical outcomes with prehabilitation. *Appl Physiol Nutr Metab* 2015; 40: 966–9.
2. Valkenet K, van de Port IG, Dronkers JJ, de Vries WR, Lindeman E, Backx FJ. The effects of preoperative exercise therapy on postoperative outcome: a systematic review. *Clin Rehabil* 2011; 25: 99–111.
3. Durrand J, Hackett R, Yates D, Danjoux G. Prehabilitation. *Perioperative Medicine – Current Controversies* 2016: 15–47.
4. Snowden CP, Prentis J, Jacques B, Anderson H, Manas D, Jones D et al. Cardiorespiratory fitness predicts mortality and hospital length of stay after major elective surgery in older people. *Ann Surg* 2013; 257: 999–1004.
5. Halder AM, Stiehler M, Günther K-P. Historie und Epidemiologie. In: *AE-Manual der Endoprothetik: Hüfte und Hüftrevision*. Berlin, Heidelberg: Springer, 2012: S. 1–20.
6. MacWilliam CH, Yood MU, Verner JJ, McCarthy BD, Ward RE. Patient-related risk factors that predict poor outcome after total hip replacement. *Health Serv Res* 1996; 31: 623–38.
7. Hermann A, Holsgaard-Larsen A, Zerahn B, Mejdahl S, Overgaard S. Preoperative progressive explosive-type resistance training is feasible and effective in patients with hip osteoarthritis scheduled for total hip arthroplasty – a randomized controlled trial. *Osteoarthritis Cartil* 2016; 24: 91–8.
8. Hoogeboom TJ, Dronkers JJ, van den Ende CHM, Oosting E, van Meeteren NLU. Preoperative therapeutic exercise in frail elderly scheduled for total hip replacement: a randomized pilot trial. *Clin Rehabil* 2010; 24: 901–10.
9. Oosting E, Jans MP, Dronkers JJ, Naber RH, Dronkers-Landman CM, Appelman-De Vries SM, et al. Preoperative Home-Based Physical Therapy Versus Usual Care to Improve Functional Health of Frail Older Adults Scheduled for Elective Total Hip Arthroplasty: A Pilot Randomized Controlled Trial. *Arch Phys Med Rehabil* 2012; 93: 610–6.
10. Bitterli R, Sieben JM, Hartmann M, de Bruin ED. Pre-surgical sensorimotor training for patients undergoing total hip replacement: a randomised controlled trial. *Int J Sports Med* 2011; 32: 725–32.
11. Ferrara PE, Rabini A, Aprile I, Maggi L, Piazzini DB, Logroscino G, et al. Effect of pre-operative physiotherapy in patients with end-stage osteoarthritis undergoing hip arthroplasty. *Clin Rehabil* 2008; 22: 977–86.
12. Gilbey HJ, Ackland TR, Wang AW, Morton AR, Troughet T, Tapper J. Exercise improves early functional recovery after total hip arthroplasty. *Clin Orthop Relat Res* 2003; 408: 193–200.
13. Gocen Z, Sen A, Karatosun V, Unver B, Karatosun V, Gunal I. The effect of preoperative physiotherapy and education on the outcome of total hip replacement: a prospective randomized controlled trial. *Clin Rehabil* 2004; 18: 353–8.

14. McGregor AH, Rylands H, Owen A, Dore CJ, Hughes SP. Does preoperative hip rehabilitation advice improve recovery and patient satisfaction? *J Arthroplasty* 2004; 19: 464–8.
15. Rooks DS, Huang J, Bierbaum BE, Bolus SA, Rubano J, Connolly CE, et al. Effect of preoperative exercise on measures of functional status in men and women undergoing total hip and knee arthroplasty. *Arthritis Rheum* 2006; 55: 700–8.
16. Villadsen A, Overgaard S, Holsgaard-Larsen A, Christensen R, Roos EM. Postoperative effects of neuromuscular exercise prior to hip or knee arthroplasty: a randomised controlled trial. *Ann Rheum Dis* 2014; 73: 1130–7.
17. Zeng R, Lin J, Wu S, Chen L, Chen S, Gao H, et al. A randomized controlled trial: preoperative home-based combined Tai Chi and Strength Training (TCST) to improve balance and aerobic capacity in patients with total hip arthroplasty (THA). *Arch Gerontol Geriatr* 2015; 60: 265–71.
18. Wang AW, Gilbey HJ, Ackland TR. Perioperative exercise programs improve early return of ambulatory function after total hip arthroplasty: a randomized, controlled trial. *Am J Phys Med Rehabil* 2002; 81: 801–6.
19. Skou ST, Odgaard A, Rasmussen JO, Roos EM. Group education and exercise is feasible in knee and hip osteoarthritis. *Dan Med J* 2012; 59: 1–5.
20. Wienand A, Cassel M, Knapp G, Reimers N, Mayer F. Arthrose. In: *Therapie und Prävention durch Sport*. Band 3. 2. Auflage. München: Urban & Fischer, 2015: S. 49–80.
21. Umehara N, Mitani S, Namba Y. Factors influencing health-related quality of life after total hip arthroplasty. *Acta Med Okayama* 2016; 70: 89–95.
22. Sampalis J, Boukas S, Liberman M, Reid T, Dupuis G. Impact of waiting time on the quality of life of patients awaiting coronary artery bypass grafting. *Can Med Assoc J* 2001; 165: 429–33.
23. Cook JW, Pierson LM, Herbert WG, Norton HJ, Fedor JM, Kiebzak GM, et al. The influence of patient strength, aerobic capacity and body composition upon outcomes after coronary artery bypass grafting. *Thorac Cardiovasc Surg* 2001; 49: 89–93.
24. Hulzebos EHJ, van Meeteren NLU, van den Buijs BJWM, de Bie RA, Brutel de la Rivière A, Helders PJM. Feasibility of preoperative inspiratory muscle training in patients undergoing coronary artery bypass surgery with a high risk of postoperative pulmonary complications: a randomized controlled pilot study. *Clin Rehabil* 2006; 20: 949–59.
25. Hulzebos EHJ, Helders PJM, Favié NJ, de Bie RA, Brutel de la Rivière A, van Meeteren NLU. Preoperative Intensive Inspiratory Muscle Training to Prevent Postoperative Pulmonary Complications in High-Risk Patients Undergoing CABG Surgery – A Randomized Clinical Trial. *JAMA* 2006; 296: 1851–7.
26. Shakouri SK, Salekzamani Y, Taghizadieh A, Sabbagh-Jadid H, Soleymani J, Sahebi L, et al. Effect of Respiratory Rehabilitation before Open Cardiac Surgery on Respiratory Function: A Randomized Clinical Trial. *J Cardiovasc Thorac Res* 2015; 7: 13–7.
27. Weiner P, Zeidan F, Zamir D, Pelled B, Waizman J, Beckerman M et al. Prophylactic Inspiratory Muscle Training in Patients Undergoing Coronary Artery Bypass Graft. *World J Surg* 1998; 22: 427–31.
28. Rosenfeldt F, Braun L, Spitzer O, Bradley S, Shepherd J, Bailey M et al. Physical conditioning and mental stress reduction – a randomised trial in patients undergoing cardiac surgery. *BMC Complement Altern Med* 2011; 11: 20.
29. Sawatzky J-A V, Kehler DS, Ready AE, Lerner N, Borek S, Lamont D et al. Prehabilitation program for elective coronary artery bypass graft surgery patients: a pilot randomized controlled study. *Clin Rehabil* 2014; 28: 648–57.
30. Arthur HM, Daniels C, McKelvie R, Hirsh J, Rush B. Effect of a Preoperative Intervention on Preoperative and Postoperative Outcomes in Low-Risk Patients Awaiting Elective. *Ann Intern Med* 2000; 133: 253–62.
31. McNeely C, Markwell S, Vassileva C. Trends in patient characteristics and outcomes of coronary artery bypass grafting in the 2000 to 2012 Medicare population. *Ann Thorac Surg* 2016; 102: 132–9.
32. Underwood MJ, Firmin RK, Juha D. Aspects of psychological and social morbidity in patients awaiting coronary artery bypass grafting. *Br Heart J* 1993; 69: 382–4.
33. Salavati M, Fallahinia G, Vardanjani AE, Rafiei H, Mousavi S, Torkamani M. Comparison Between Effects of Home Based Cardiac Rehabilitation Programs Versus Usual Care on the Patients' Health Related Quality of Life After Coronary Artery Bypass Graft. *Glob J Health Sci* 2016; 8: 196–202.
34. Hulzebos EHJ, Smit Y, Helders PJM, van Meeteren NLU. Preoperative physical therapy for elective cardiac surgery patients. *Cochrane Database Syst Rev* 2012; 11: CD010118.
35. Chang JI, Lam V, Patel MI. Preoperative pelvic floor muscle exercise and postprostatectomy incontinence: a systematic review and meta-analysis. *Eur Urol* 2016; 69: 460–7.
36. MacDonald R, Fink HA, Huckabay C, Monga M, Wilt TJ. Pelvic floor muscle training to improve urinary incontinence after radical prostatectomy: A systematic review of effectiveness. *BJU Int* 2007; 100: 76–81.
37. Zopf EM, Baumann FT. Prostatakrebs. In: *Sport und körperliche Aktivität in der Onkologie*. Berlin, Heidelberg: Springer, 2012: S. 179–88.
38. Burgio KL, Goode PS, Urban DA, Umlauf MG, Locher JL, Bueschen A, et al. Preoperative biofeedback assisted behavioral training to decrease post-prostatectomy incontinence: a randomized, controlled trial. *J Urol* 2006; 175: 196–201.
39. Dijkstra-Eshuis J, van den Bos TWL, Splinter R, Bevers RFM, Zonneveld WCG, Putter H et al. Effect of preoperative pelvic floor muscle therapy with biofeedback versus standard care on stress urinary incontinence and quality of life in men undergoing laparoscopic radical prostatectomy: a randomised control trial. *Neurourol Urodyn* 2015; 34: 144–50.

40. Geraerts I, van Poppel H, Devoogdt N, Joniau S, van Cleynebreugel B, de Groef A, et al. Influence of preoperative and postoperative pelvic floor muscle training (PFMT) compared with postoperative PFMT on urinary incontinence after radical prostatectomy: A randomized controlled trial. *Eur Urol* 2013; 64: 766–72.
41. Laurienzo CE, Sacomani CAR, Rodrigues TR, Zequi S de CS de C, Guimarães GC, Lopes A. Results of preoperative electrical stimulation of pelvic floor muscles in the continence status following radical retropubic prostatectomy. *Int Brazilian J Urol* 2013; 39: 182–8.
42. Centemero A, Rigatti L, Giraudo D, Lazzeri M, Lughezzani G, Zugna D et al. Preoperative pelvic floor muscle exercise for early continence after radical prostatectomy: a randomised controlled study. *Eur Urol* 2010; 57: 1039–43.
43. Sueppel C, Kreder K, See W. Improved Continence Outcomes With Preoperative Pelvic Floor Muscle Strengthening Exercises. *Urol Nurs* 2001; 21: 201–10.
44. Walsh PC, Marschke P, Ricker D, Burnett AL. Patient-reported urinary continence and sexual function after anatomic radical prostatectomy. *Urology* 2000; 55: 58–61.
45. Hirakawa T, Suzuki S, Kato K, Gotoh M, Yoshikawa Y. Randomized controlled trial of pelvic floor muscle training with or without biofeedback for urinary incontinence. *Int Urogynecol J* 2013; 24: 1347–54.

Literatur zu Kapitel 9 Koronare Herzkrankheit

1. Statistisches Bundesamt 2016 <https://www.destatis.de/DE/ZahlenFakten/GesellschaftStaat/Gesundheit/Todesursachen/Todesursachen.html;jsessionid=DADDFCFDDE.cae>.
2. Nationale Versorgungsleitlinie Chronische KHK, Langfassung. 4. Auflage 2016 www.awmf.org/uploads/tx_szleitlinien/nvl-004I_S3_KHK_2016-02.pdf
3. Piepoli MF, Hoes AW, Agewall S, Albus C, Brotons C et al. 2016 European Guidelines on cardiovascular disease prevention in clinical practice: The Sixth Joint Task Force of the European Society of Cardiology and Other Societies on Cardiovascular Disease Prevention in Clinical Practice. *Eur Heart J* 2016; 37: 36: 2768–2801.
4. Global status report on noncommunicable diseases 2014. 1. Chronic Disease – prevention and control. 2. Chronic Disease – epidemiology. 3. Chronic Disease – mortality. 4. Cost of Illness. 5. Delivery of Health Care. I. World Health Organization. ISBN 978 92 4 156485 4. WHO, Genf 2014. (www.who.int)
5. Bjarnason-Wehrens B, Mai M, Knapp G. Koronare Herzkrankheit. In: Mooren F, Knapp G, Reimers C. D. (Hrsg). Prävention und Therapie durch Sport. Band 4 Innere Medizin/Pädiatrie. Urban & Fischer, München 2016, S. 169–255.
6. Sattelmair JR, Perma J, Ding EL, Kohl HW, Haskell W, Lee IM. Dose-response between physical activity and risk of coronary heart disease: a meta-analysis. *Circulation* 2011; 124: 789–95.
7. Bjarnason-Wehrens B, Schulz O, Gielen S, Halle M, Dürsch M et al. Leitlinie körperliche Aktivität zur Sekundärprävention und Therapie kardiovaskulärer Erkrankungen. *Clin Res Cardiol* 2009; 7 Suppl. 3: 1–44.
8. Vanhees L, Rauch B, Piepoli M, van Buuren F, Takken T et al. Importance of characteristics and modalities of physical activity and exercise in the management of cardiovascular health in individuals with cardiovascular disease (Part III). *Eur J Prev Cardiol* 2012; 19: 6:1333–56.
9. Piepoli MF, Corrà U, Benzer W, Bjarnason-Wehrens B, Dendale P et al. Secondary prevention through cardiac rehabilitation: from knowledge to implementation. A position paper from the cardiac rehabilitation section of the European Association of Cardiovascular Prevention and Rehabilitation. *Eur J Cardiovasc Prev Rehabil* 2010; 17: 1–17.
10. Piepoli MF, Corrà U, Dendale P, Frederix I, Prescott E et al. Challenges in secondary prevention after acute myocardial infarction: A call for action. *Eur J Prev Cardiol* 2016 Sep 6. pii: 23:18:1994-2006.
11. Anderson L, Oldridge N, Thompson DR, Zwisler AD, Rees K et al. Exercise-Based Cardiac Rehabilitation for Coronary Heart Disease: Cochrane Systematic Review and Meta-Analysis. *J Am Coll Cardiol* 2016 Jan 5; 67: 1–12. doi: 10.1016/j.jacc.2015.10.044.
12. Rauch B, Davos CH, Doherty P, Saure D, Metzendorf MI et al. The prognostic effect of cardiac rehabilitation in the era of acute revascularisation and statin therapy: A systematic review and meta-analysis of randomized and non-randomized studies – The Cardiac Rehabilitation Outcome Study (CROS). *Eur J Prev Cardiol* 2016 Oct 24. pii: 23:18:1914-39.
13. Vanhees L, De Sutter J, Gelada SN, Doyle F, Prescott E et al. Importance of characteristics and modalities of physical activity and exercise in defining the benefits to cardiovascular health within the general population: recommendations from the EACPR (Part I). *Eur J Prev Cardiol* 2012; 19: 670–86.
14. Ruiz JR, Sui X, Lobelo F, Morrow JR Jr, Jackson AW et al. Association between muscular strength and mortality in men: prospective cohort study. *BMJ* 2008; 337: 439.
15. Zhao G, Li C, Ford ES, et al. Fulton JE, Carlson SA, et al. Leisure-time aerobic physical activity, muscle-strengthening activity and mortality risks among US adults: the NHANES linked mortality study. *Br J Sports Med* 2014; 48: 244–9.
16. Arturo EG, Lee DC, Lavie CJ, España-Romero V, Sui X, et al. Effects of muscular strength on cardiovascular risk factors and prognosis. *J Cardiopulm Rehabil Prev* 2012; 32: 351–8.

17. Marzolini S, Oh PI, Brooks D. Effect of combined aerobic and resistance training versus aerobic training alone in individuals with coronary artery disease: a meta-analysis. *Eur J Prev Cardiol* 2012; 19: 81–94.
18. Yamamoto S, Hotta K, Ota E, Mori R, Matsunaga A. Effects of resistance training on muscle strength, exercise capacity, and mobility in middle-aged and elderly patients with coronary artery disease: A meta-analysis. *J Cardiol* 2016; 68: 125–34.
19. Buchheit M, Laursen PB. High-intensity interval training, solutions to the programming puzzle: Part I: cardiopulmonary emphasis. *Sports Med* 2013; 43: 313–38.
20. Weston KS, Wisløff U, Coombes JS. High-intensity interval training in patients with lifestyle-induced cardiometabolic disease: a systematic review and meta-analysis. *Br J Sports Med* 2014; 48: 1227–34.
21. Milanović Z, Sporiš G, Weston M. Effectiveness of High-Intensity Interval Training (HIT) and Continuous Endurance Training for VO2max Improvements: A Systematic Review and Meta-Analysis of Controlled Trials. *Sports Med* 2015; 45: 1469–81.
22. Jelleyman C, Yates T, O'Donovan G, Gray LJ, King JA et al. The effects of high-intensity interval training on glucose regulation and insulin resistance: a meta-analysis. *Obes Rev* 2015; 16: 942–61.
23. Pattyn N, Coeckelberghs E, Buys R, Cornelissen VA, Vanhees L. Aerobic interval training vs. moderate continuous training in coronary artery disease patients: a systematic review and meta-analysis. *Sports Med* 2014; 44: 687–700.
24. Elliott AD, Rajopadhyaya K, Bentley DJ, Beltrame JF, Aromataris EC. Interval Training Versus Continuous Exercise in Patients with Coronary Artery Disease A Meta-Analysis. *Heart, Lung and Circulation* 2015; 24: 149–57.
25. Conraads VM, Pattyn N, De Maeyer C, Beckers PJ, Coeckelberghs E et al. Aerobic interval training and continuous training equally improve aerobic exercise capacity in patients with coronary artery disease; The SAINTEX-CAD study. *Intern J Cardiology* 2015; 179: 203–10.
26. Pattyn N, Vanhees L, Cornelissen VA, Coeckelberghs E, De Maeyer C et al. The long-term effects of a randomized trial comparing aerobic interval versus continuous training in coronary artery disease patients: 1-year data from the SAINTEX-CAD study. *Eur J Prev Cardiol* 2016; Doi: 2047487316631200.
27. Borjesson M, Urhausen A, Koudi E, Dugmore D, Sharma S et al. Cardiovascular evaluation of middle-aged/senior individuals engaged in leisure-time sport activities: position stand from the sections of exercise physiology and sports cardiology of the European Association of Cardiovascular Prevention and Rehabilitation. *Eur J Cardiovasc Prev Rehabil* 2011; 18: 446–58.
28. Bjarnason-Wehrens B, Halle M. Exercise Training in Cardiac Rehabilitation. In: Niebauer J. (Ed) *Cardia Rehabilitation Manual* (2. Ed.), Springer, London 2016, S. 89–120.
29. Kraemer WJ, Ratamess NA, Fry AC, French DN. Strength testing: development and evaluation of methodology. In: Maud PJ, Foster C, eds. *Physiological assessment of human fitness*. 2nd ed. Champaign, IL: Human Kinetics; 2006:19–150.

Literatur zu Kapitel 10 Körperliches Training bei chronischer Herzinsuffizienz

1. Ponikowski P, Voors AA, Anker SD, Bueno H, Cleland JGF, Coats AJS et al. 2016 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure. *Eur Heart J* 2015; ehv128.
2. Störk S, Frantz S, Bauersachs J, Ertl G, Angermann CE. Primary diagnosis of heart failure in ambulatory and hospitalized patients. *Dtsch Med Wochenschr* 2008; 133: 636–41.
3. Coats AJS. What causes the symptoms of heart failure? *Heart* 2001; 86: 574–8.
4. Packer M. Abnormalities of diastolic function as a potential cause of exercise intolerance in chronic heart failure. *Circulation*. Februar 1990;81(2 Suppl):III78–86.
5. Sandri M, Kozarez I, Adams V, Mangner N, Höllriegel R, Erbs S et al. Age-related effects of exercise training on diastolic function in heart failure with reduced ejection fraction: the Leipzig Exercise Intervention in Chronic Heart Failure and Aging (LEICA) Diastolic Dysfunction Study. *Eur Heart J* 2012; 33: 1758–68.
6. Gielen S, Adams V, Möbius-Winkler S, Linke A, Erbs S, Yu J et al. Anti-inflammatory effects of exercise training in the skeletal muscle of patients with chronic heart failure. *J Am Coll Cardiol* 2003; 42: 861–8.
7. Gielen S, Adams V, Linke A, Erbs S, Möbius-Winkler S, Schubert A et al. Exercise training in chronic heart failure: correlation between reduced local inflammation and improved oxidative capacity in the skeletal muscle. *Eur J Cardiovasc Prev Rehabil* 2005; 12: 393–400.
8. Hambrecht R, Fiehn E, Yu J, Niebauer J, Weigl C, Hilbrich L et al. Effects of endurance training on mitochondrial ultrastructure and fiber type distribution in skeletal muscle of patients with stable chronic heart failure. *J Am Coll Cardiol* 1997; 29: 1067–73.
9. Gielen S, Sandri M, Kozarez I, Kratzsch J, Teupser D, Thiery J et al. Exercise training attenuates MuRF-1 expression in the skeletal muscle of patients with chronic heart failure independent of age: the randomized Leipzig Exercise Intervention in Chronic Heart Failure and Aging catabolism study. *Circulation* 2012; 125: 2716–27.

10. Anker SD, Coats AJ. Cardiac cachexia: a syndrome with impaired survival and immune and neuroendocrine activation. *Chest* 1999; 115: 836–47.
11. Strassburg S, Springer J, Anker SD. Muscle wasting in cardiac cachexia. *Int J Biochem Biol* 2005; 37: 1938–47.
12. Coats AJ, Clark AL, Piepoli M, Volterrani M, Poole-Wilson PA. Symptoms and quality of life in heart failure: the muscle hypothesis. *Br Heart J* 1994; 72 (2 Suppl): S36–9.
13. Coats AJ. The muscle hypothesis of chronic heart failure. *J Mol Cell Cardiol* 1996; 28: 2255–62.
14. Piepoli MF, Guazzi M, Boriani G, Ciccoira M, Corrà U, Dalla Libera L et al. Exercise intolerance in chronic heart failure: mechanisms and therapies. Part I. *Eur J Cardiovasc Prev Rehabil Off J Eur Soc Cardiol Work Groups Epidemiol Prev Card Rehabil Exerc Physiol* 2010; 17: 637–42.
15. Coats AJS, Adamopoulos S, Meyer TE, Conway J, Sleight P. Effects of physical training in chronic heart failure. *The Lancet* 1990; 335: 63–6.
16. Taylor RS, Sagar VA, Davies EJ, Briscoe S, Coats AJS, Dalal H et al. Exercise-based rehabilitation for heart failure. *Cochrane Database Syst Rev* 2014; 4: CD003331.
17. Hambrecht R, Gielen S, Linke A, Fiehn E, Yu J, Walther C et al. Effects of exercise training on left ventricular function and peripheral resistance in patients with chronic heart failure: A randomized trial. *JAMA J Am Med Assoc* 2000; 283: 3095–101.
18. Smart N, Haluska B, Jeffriess L, Marwick TH. Exercise training in systolic and diastolic dysfunction: effects on cardiac function, functional capacity, and quality of life. *Am Heart J* 2007; 153: 530–6.
19. Kubo SH, Rector TS, Williams RE, Heifritz SM, Bank AJ. Endothelium dependent vasodilation is attenuated in patients with heart failure. *Circulation* 1994; 84: 1589–96.
20. Ismail H, McFarlane JR, Nojournian AH, Dieberg G, Smart NA. Clinical outcomes and cardiovascular responses to different exercise training intensities in patients with heart failure: a systematic review and meta-analysis. *JACC Heart Fail* 2013; 1: 514–22.
21. Piepoli MF, Davos C, Francis DP, Coats AJ. Exercise training meta-analysis of trials in patients with chronic heart failure (ExTraMATCH). *BMJ* 2004; 328: 189.
22. O'Connor CM, Whellan DJ, Lee KL, Keteyian SJ, Cooper LS, Ellis SJ et al. Efficacy and safety of exercise training in patients with chronic heart failure: HF-ACTION randomized controlled trial. *JAMA J Am Med Assoc* 2009; 301: 1439–50.
23. Keteyian SJ, Leifer ES, Houston-Miller N, Kraus WE, Brawner CA, O'Connor CM et al. Relation between volume of exercise and clinical outcomes in patients with heart failure. *J Am Coll Cardiol* 2012; 60: 1899–905.
24. Wisloff U, Støylen A, Loennechen JP, Bruvold M, Rognmo O, Haram PM et al. Superior cardiovascular effect of aerobic interval training versus moderate continuous training in heart failure patients: a randomized study. *Circulation* 2007; 115: 3086–94.
25. Ellingsen Ø, Halle M, Conraads V, Støylen A, Dalen H, Delagardelle C et al. High-Intensity Interval Training in Patients With Heart Failure With Reduced Ejection Fraction. *Circulation* 2017; 135: 839–49.
26. Bjarnason-Wehrens B, Mayer-Berger W, Meister ER, Baum K, Hambrecht R, Gielen S. Resistance exercise in cardiac rehabilitation-recommendations of the German Federation for Cardiovascular Prevention and Rehabilitation. *Z Kardiol* 2004; 93: 357–70.
27. Pu CT, Johnson MT, Forman DE, Hausdorff JM, Roubenoff R, Foldvari M et al. Randomized trial of progressive resistance training to counteract the myopathy of chronic heart failure. *J Appl Physiol* 2001; 90: 2341–50.
28. Scheinowitz M, Harpaz D. Safety of cardiac rehabilitation in a medically supervised, community-based program. *Cardiology* 2005; 103: 113–7.
29. Mittleman MA. Angina in patients with an active lifestyle. *Eur Heart J* 1996; 17 Suppl G: 30–5.
30. Dahabreh IJ, Paulus JK. Association of episodic physical and sexual activity with triggering of acute cardiac events: systematic review and meta-analysis. *JAMA* 2011; 305: 1225–33.
31. Pavy B, Iliou MC, Meurin P, Tabet J-Y, Corone S, Functional Evaluation and Cardiac Rehabilitation Working Group of the French Society of Cardiology. Safety of exercise training for cardiac patients: results of the French registry of complications during cardiac rehabilitation. *Arch Intern Med* 2006; 166: 2329–34.
32. Piepoli MF, Conraads V, Corrà U, Dickstein K, Francis DP, Jaarsma T et al. Exercise training in heart failure: from theory to practice. A consensus document of the Heart Failure Association and the European Association for Cardiovascular Prevention and Rehabilitation. *Eur J Heart Fail* 2011; 13: 347–57.
33. Edelmann F, Gelbrich G, Dungen H-D, Fröhling S, Wachter R, Stahrenberg R et al. Exercise training improves exercise capacity and diastolic function in patients with heart failure with preserved ejection fraction: results of the Ex-DHF (Exercise training in Diastolic Heart Failure) pilot study. *J Am Coll Cardiol* 2011; 58: 1780–91.
34. Nolte K, Herrmann-Lingen C, Wachter R, Gelbrich G, Dungen H-D, Duvinage A et al. Effects of exercise training on different quality of life dimensions in heart failure with preserved ejection fraction: the Ex-DHF-P trial. *Eur J Prev Cardiol* 2015; 22: 582–93.
35. Pandey A, Parashar A, Kumbhani DJ, Agarwal S, Garg J, Kitzman D et al. Exercise training in patients with heart failure and preserved ejection fraction: meta-analysis of randomized control trials. *Circ Heart Fail* 2015; 8: 33–40.

36. Vanhees L, Schepers D, Heidebuchel H, Defoor J, Fagard R. Exercise performance and training in patients with implantable cardioverter-defibrillators and coronary heart disease. *Am J Cardiol* 2001; 87: 712–5.
37. Vanhees L, Kornaat M, Defoor J, Aufdemkampe G, Schepers D, Stevens A et al. Effect of exercise training in patients with an implantable cardioverter defibrillator. *Eur Heart J* 2004; 25: 1120–6.
38. Moe GW, Ezekowitz JA, O'Meara E, Lepage S, Howlett JG, Fremes S et al. The 2014 Canadian Cardiovascular Society Heart Failure Management Guidelines Focus Update: anemia, biomarkers, and recent therapeutic trial implications. *Can J Cardiol* 2015; 31: 3–16.
39. Kitzman DW, Brubaker PH, Morgan TM, Stewart KP, Little WC. Exercise training in older patients with heart failure and preserved ejection fraction: a randomized, controlled, single-blind trial. *Circ Heart Fail* 2010; 3: 659–67.
40. Alves AJ, Ribeiro F, Goldhammer E, Rivlin Y, Rosenschein U, Viana JL et al. Exercise training improves diastolic function in heart failure patients. *Med Sci Sports Exerc* 2012; 44: 776–85.
41. Smart NA, Haluska B, Jeffries L, Leung D. Exercise training in heart failure with preserved systolic function: a randomized controlled trial of the effects on cardiac function and functional capacity. *Congest Heart Fail Greenwich Conn* 2012; 18: 295–301.
42. Isaksen K, Morken IM, Munk PS, Larsen AI. Exercise training and cardiac rehabilitation in patients with implantable cardioverter defibrillators: a review of current literature focusing on safety, effects of exercise training, and the psychological impact of programme participation. *Eur J Prev Cardiol* 2012; 19: 804–12.
43. Piepoli MF, Corrà U, Adamopoulos S, Benzer W, Bjarnason-Wehrens B, Cupples M et al. Secondary prevention in the clinical management of patients with cardiovascular diseases. Core components, standards and outcome measures for referral and delivery: a policy statement from the cardiac rehabilitation section of the European Association for Cardiovascular Prevention & Rehabilitation. Endorsed by the Committee for Practice Guidelines of the European Society of Cardiology. *Eur J Prev Cardiol* 2014; 21: 664–81.
44. Piepoli MF, Corrà U, Benzer W, Bjarnason-Wehrens B, Dendale P, Gaita D et al. Secondary prevention through cardiac rehabilitation: from knowledge to implementation. A position paper from the Cardiac Rehabilitation Section of the European Association of Cardiovascular Prevention and Rehabilitation. *Eur J Cardiovasc Prev Rehabil Off J Eur Soc Cardiol Work Groups Epidemiol Prev Card Rehabil Exerc Physiol* 2010; 17: 1–17.
45. Niebauer J, Mayr K, Tschentscher M, Pokan R, Benzer W. Outpatient cardiac rehabilitation: the Austrian model. *Eur J Prev Cardiol* 2013; 20: 468–79.

Literatur zu Kapitel 11 Arterielle Hypertonie

1. Gilbert N, Hager A. Therapie der arteriellen Hypertonie bei Kindern und Jugendlichen. Aktualisierte Therapieempfehlungen. *Arzneimitteltherapie* 2014; 32: 234–44.
2. James PA, Oparil S, Carter BL, Cushman WC, Dennison-Himmelfarb C et al. 2014 evidence-based guideline for the management of high blood pressure in adults: report from the panel members appointed to the Eighth Joint National Committee (JNC 8). *JAMA* 2014; 311: 507–20.
3. Deutsche Gesellschaft für Kardiologie – Herz- und Kreislaufforschung e.V., Deutsche Hochdruckliga e.V., Deutsche Gesellschaft für Hypertonie und Prävention. ESC POCKET GUIDELINES. Leitlinien für das Management der arteriellen Hypertonie, 2013.
4. Janhsen K, Strube H, Starker A: Hypertonie. Gesundheitsberichterstattung des Bundes. Heft 43. Robert-Koch-Institut, Statistisches Bundesamt, 2008.
5. Kitchen TA. Hypertensive vascular disease. In: Longo DL, Fauci AS, Kasper DL, Hauser SL, Jameson JL, Loscalzo J (eds.) *Harrison's principles of internal medicine*. Vol II, New York, Chicago, San Francisco, Lisbon, London, Madrid, Mexico City, Milan, New Delhi, San Juan, Seoul, Singapore, Sydney, Toronto: McGraw-Hill, 2012, pp. 2042–2059.
6. Rickenbacher P. Update Arterielle Hypertonie 2015. *Praxis* 2015; 104: 461–5.
7. Reimers CD, Knapp G (2016) Arterielle Hypertonie. In: Mooren F, Knapp G, Reimers CD (Hrsg.) *Prävention und Therapie durch Sport*. Band 4. Innere Medizin/Pädiatrie. 2. Aufl., München: Urban & Fischer, S. 121–68.
8. Predel HG: Bluthochdruck und Sport. *Dt Zeitschr Sportmed* 2007; 58: 328–33.
9. Owen A, Wiles J, Swaine I. Effect of isometric exercise on resting blood pressure: a meta analysis. *J Hum Hypertens* 2010; 24: 796–800.
10. Whelton SP, Chin A, Xin X, He J. Effect of aerobic exercise on blood pressure: a meta-analysis of randomized, controlled trials. *Ann Intern Med* 2002; 136: 493–503.
11. Cornelissen VA, Buys R, Smart NA. Endurance exercise beneficially affects ambulatory blood pressure: a systematic review and meta-analysis. *J Hypertens* 2013; 31: 639–48.
12. Cornelissen VA, Fagard RH. Effects of endurance training on blood pressure, blood pressure-regulating mechanisms, and cardiovascular risk factors. *Hypertension* 2005; 46: 667–75.

13. Cornelissen VA, Fagard RH, Coeckelberghs E, Vanhees L. Impact of resistance training on blood pressure and other cardiovascular risk factors: a meta-analysis of randomized, controlled trials. *Hypertension* 2011; 58: 950–8.
14. Lee MS, Pittler MH, Guo R, Ernst E. Qigong for hypertension: a systematic review of randomized clinical trials. *J Hypertens* 2007; 25: 1525–32.
15. Gaciong Z, Sinski M, Lewandowski J. Blood pressure control and primary prevention of stroke: Summary of the recent clinical trial data and meta-analyses. *Curr Hypertens Rep* 2013; 15: 559–74.
16. Cesa CC, Sbruzzi G, Ribeiro RA, Baarbiero SM, de Oliveira Petkoicz R et al. Physical activity and cardiovascular risk factors in children: meta-analysis of randomized clinical trials. *Prev Med* 2014; 69: 54–62.
17. Gkaliagkousi E, Gavrilaki E, Douma S. Effects of acute and chronic exercise in patients with essential hypertension: benefits and risks. *Am J Hypertens* 2015; 28: 429–39.
18. Deutsche Liga zur Bekämpfung des hohen Blutdruckes e. V. Deutsche Hypertonie Gesellschaft. *Hypertonie und Sport*. 3. Aufl., 1998.
19. McDonnell BJ, Maki-Petaja KM, Munnery M, Wilkinson YIB, Cockcroft JR, McEnery CM. Habitual exercise and blood pressure: age dependency and underlying mechanisms. *Am J Hypertens* 2013; 26: 334–41.

Literatur zu Kapitel 12 Periphere arterielle Verschlusskrankheit

1. Hirsch AT, Haskal ZJ, Hertzner NR, Bakal CW, Creager MA et al. ACC/AHA 2005 Practice Guidelines for the management of patients with peripheral arterial disease (lower extremity, renal, mesenteric, and abdominal aortic): a collaborative report from the American Association for Vascular Surgery/Society for Vascular Surgery, Society for Cardiovascular Angiography and Interventions, Society for Vascular Medicine and Biology, Society of Interventional Radiology, and the ACC/AHA Task Force on Practice Guidelines (Writing Committee to Develop Guidelines for the Management of Patients With Peripheral Arterial Disease): endorsed by the American Association of Cardiovascular and Pulmonary Rehabilitation; National Heart, Lung, and Blood Institute; Society for Vascular Nursing; TransAtlantic Inter-Society Consensus; and Vascular Disease Foundation. *Circulation* 2006; 113: e463–654.
2. Rooke TW, Hirsch AT, Misra S, Sidawy AN, Beckman JA et al. 2011 ACCF/AHA focused update of the guideline for the management of patients with peripheral artery disease (updating the 2005 guideline): a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines: developed in collaboration with the Society for Cardiovascular Angiography and Interventions, Society of Interventional Radiology, Society for Vascular Medicine, and Society for Vascular Surgery. *Catheter Cardiovasc Interv* 2012; 79: 501–31.
3. Schmidt-Trucksäss A, Schmid A, Brunner C, Scherer N, Zäch G et al. Arterial properties of the carotid and femoral artery in endurance-trained and paraplegic subjects. *J Appl Physiol* 2000; 89: 1956–63.
4. Stewart KJ, Hiatt WR, Regensteiner JG, Hirsch AT. Exercise training for claudication. *N Engl J Med* 2002; 347: 1941–51.
5. Treat-Jacobson D, Bronas UG, Leon AS. Efficacy of arm-ergometry versus treadmill exercise training to improve walking distance in patients with claudication. *Vasc Med* 2009; 14: 203–13.
6. Thijssen DH, Green DJ, Steendijk S, Hopman MT. Sympathetic vasomotor control does not explain the change in femoral artery shear rate pattern during arm-crank exercise. *Am J Physiol Heart Circ Physiol* 2009; 296: H180–5.
7. Lane R, Ellis B, Watson L, Leng GC. Exercise for intermittent claudication. *Cochrane Database Syst Rev* 2014; 7: CD000990.
8. Watson L, Ellis B, Leng GC. Exercise for intermittent claudication. *Cochrane Database Syst Rev* 2008:CD000990.
9. Gardner AW, Killewich LA, Montgomery PS, Katzel LI. Response to exercise rehabilitation in smoking and nonsmoking patients with intermittent claudication. *J Vasc Surg* 2004; 39: 531–8.
10. McDermott MM, Ades P, Guralnik JM, Dyer A, Ferrucci L et al. Treadmill exercise and resistance training in patients with peripheral arterial disease with and without intermittent claudication: a randomized controlled trial. *JAMA* 2009; 301: 165–74.
11. Ritti-Dias RM, Wolosker N, de Moraes Forjaz CL, Carvalho CR, Cucato GG et al. Strength training increases walking tolerance in intermittent claudication patients: randomized trial. *J Vasc Surg* 2010; 51: 89–95.
12. Tuner SL, Easton C, Wilson J, Byrne DS, Rogers P et al. Cardiopulmonary responses to treadmill and cycle ergometry exercise in patients with peripheral vascular disease. *J Vasc Surg* 2008; 47: 123–30.
13. Sieminski DJ, Gardner AW. The relationship between free-living daily physical activity and the severity of peripheral arterial occlusive disease. *Vasc Med* 1997; 2: 286–91.
14. Nicolai SP, Hendriks EJ, Prins MH, Teijink JA. Optimizing supervised exercise therapy for patients with intermittent claudication. *J Vasc Surg* 2010; 52: 1226–33.
15. Gardner AW, Poehlman ET. Exercise rehabilitation programs for the treatment of claudication pain. A meta-analysis. *JAMA* 1995; 274: 975–80.

16. Parmenter BJ, Raymond J, Dinnen P, Singh MA. A systematic review of randomized controlled trials: Walking versus alternative exercise prescription as treatment for intermittent claudication. *Atherosclerosis* 2011; 218: 1–12.
17. Fokkenrood HJ, Bendermacher BL, Lauret GJ, Willigendael EM, Prins MH, Teijink JA. Supervised exercise therapy versus non-supervised exercise therapy for intermittent claudication. *Cochrane Database Syst Rev* 2013; 8: CD005263.

Literatur zu Kapitel 13 Diabetes mellitus

1. Kerner W, Brückel J. Definition, Klassifikation und Diagnostik des Diabetes mellitus. *Diabetologie* 2013; 8: S104–S107.
2. Mooren FC. Diabetes mellitus Typ 2 und Sport – Blutzuckerkontrolle durch Ausdauer- und Krafttraining. *Kliniker* 2013; 42: 406–410.
3. Kemmer FW, Halle M, Stumvoll M, Thurm U, Zimmer P. Diabetes, Sport und Bewegung. *Diabetologie* 2012; 7: S170–S173.
4. Colberg SR, Sigal RJ, Fernhall B, Regensteiner JG, Blissmer BJ, et al. American College of Sports Medicine; American Diabetes Association. Exercise and type 2 diabetes: the American College of Sports Medicine and the American Diabetes Association: joint position statement. *Diabetes Care* 2010 Dec; 33: e147–67.
5. Helmrich SP, Ragland DR, Leung RW, Paffenbarger RS Jr. Physical activity and reduced occurrence of non-insulin-dependent diabetes mellitus. *N Engl J Med* 1991 Jul 18; 325: 147–52.
6. Grøntved A, Pan A, Mekary RA, Stampfer M, Willett WC, et al. Muscle-strengthening and conditioning activities and risk of type 2 diabetes: a prospective study in two cohorts of US women. *PLoS Med* 2014 Jan; 11: e1001587.
7. Tuomilehto J, Lindström J, Eriksson JG, Valle TT, Hämäläinen H, et al; Finnish Diabetes Prevention Study Group. Prevention of type 2 diabetes mellitus by changes in lifestyle among subjects with impaired glucose tolerance. *N Engl J Med* 2001 May 3; 344: 1343–50.
8. Knowler WC, Barrett-Connor E, Fowler SE, Hamman RF, Lachin JM, Walker EA, Nathan DM; Diabetes Prevention Program Research Group. Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *N Engl J Med* 2002 Feb 7; 346: 393–403.
9. Orozco LJ, Buchleitner AM, Gimenez-Perez G, Roqué I Figuls M, Richter B, Mauricio D. Exercise or exercise and diet for preventing type 2 diabetes mellitus. *Cochrane Database Syst Rev* 2008 Jul 16:CD003054.
10. Sigal RJ, Kenny GP, Boulé NG, Wells GA, Prud'homme D, Fortier M, Reid RD, Tulloch H, Coyle D, Phillips P, Jennings A, Jaffey J. Effects of aerobic training, resistance training, or both on glycemic control in type 2 diabetes: a randomized trial. *Ann Intern Med* 2007; 147: 357–69.
11. Church TS, Blair SN, Cocroham S, Johannsen N, Johnson W, Kramer K, Mikus CR, Myers V, Nauta M, Rodarte RQ, Sparks L, Thompson A, Earnest CP. Effects of aerobic and resistance training on hemoglobin A1c levels in patients with type 2 diabetes: a randomized controlled trial. *JAMA* 2010; 304: 2253–62.
12. Boulé NG, Haddad E, Kenny GP, Wells GA, Sigal RJ. Effects of exercise on glycemic control and body mass in type 2 diabetes mellitus: a meta-analysis of controlled clinical trials. *JAMA* 2001 Sep 12; 286: 1218–27.
13. Umpierre D, Ribeiro PA, Kramer CK, Leitão CB, Zucatti AT, et al. Physical activity advice only or structured exercise training and association with HbA1c levels in type 2 diabetes: a systematic review and meta-analysis. *JAMA*. 2011 May 4; 305: 1790–9.
14. Balducci S, Sacchetti M, Haxhi J, Orlando G, D'Errico V, et al. Physical exercise as therapy for type 2 diabetes mellitus. *Diabetes Metab Res Rev* 2014 Mar; 30 Suppl 1: 13–23.
15. Jorge ML1, de Oliveira VN, Resende NM, Paraiso LF, Calixto A, et al. The effects of aerobic, resistance, and combined exercise on metabolic control, inflammatory markers, adipocytokines, and muscle insulin signaling in patients with type 2 diabetes mellitus. *Metabolism* 2011 Sep; 60: 1244–52. doi: 10.1016/j.metabol.2011.01.006. Epub 2011 Mar 4.
16. Kadooglou NP, Fotiadis G, Kapelouzou A, Kostakis A, Liapis CD, Vrabas IS. The differential anti-inflammatory effects of exercise modalities and their association with early carotid atherosclerosis progression in patients with type 2 diabetes. *Diabet Med* 2013 Feb; 30: e41–50.

Literatur zu Kapitel 14 Übergewicht und Adipositas

1. Berg A, Bischoff SC, Colombo-Benkmann M, et al. Interdisziplinäre Leitlinie der Qualität S3 zur „Prävention und Therapie der Adipositas“. 2. ed. Martinsried: Deutsche Adipositas Gesellschaft, 2011: 1.
2. Obesity: preventing and managing the global epidemic. Report of a WHO consultation. *World Health Organ Tech Rep Ser* 2000; 894: i-253.
3. Die Nationale Verzehrsstudie II. Max Rubner-Institut. 1–143. 2008. Bundesforschungsinstitut für Ernährung und Lebensmittel.

4. Mensink GB. Physical activity. *Gesundheitswesen* 1999; 61 Spec No: S126–S131.
5. Löllgen H. Primärprävention kardialer Erkrankungen: Stellenwert der körperlichen Aktivität. *Deutsch Ärzteblatt* 2003; 100: A-987–A-996.
6. Mayer F, Gollhofer A, Berg A. Krafttraining mit Älteren und chronisch Kranken. *Deutsch Z Sportmed* 2003; 54.
7. Berg A, Jakob E, Lehmann M, Dickhuth HH, Huber G, Keul J. Aktuelle Aspekte der modernen Ergometrie. *Pneumologie* 1990; 44: 2–13.
8. Berg A, Halle M, Franz I, Keul J. Physical activity and lipoprotein metabolism: Epidemiological evidence and clinical trials. *Eur J Med Res* 1997; 2: 259–64.
9. Hurley BF, Nemeth PM, Martin WH, Hagberg JM, Dalsky GP, Holloszy JO. Muscle triglyceride utilisation during exercise: effect of training. *J Appl Physiol* 1986; 60: 562–7.
10. Romijn JA, Coyle EF, Sidossis LS, Rosenblatt J, Wolfe RR. Substrate metabolism during different exercise intensities in endurance-trained women. *J Appl Physiol* 2000; 88: 1707–14.
11. Bijnen FC, Feskens EJ, Caspersen CJ, et al. Physical activity and cardiovascular risk factors among elderly men in Finland, Italy, and the Netherlands. *Am J Epidemiol* 1996; 143: 553–61.
12. Bouchard C, An P, Rice T, et al. Familial aggregation of VO(2max) response to exercise training: results from the HERITAGE Family Study. *J Appl Physiol* 1999; 87: 1003–8.
13. Berg A, Halle M. Körperliche Aktivität und kardiovaskuläre Mortalität: Von der Epidemiologie zur medizinischen Praxis. *Medizinische Welt* 1999; 50: 359–62.
14. Donnelly JE, Blair SN, Jakicic JM, Manore MM, Rankin JW, Smith BK. American College of Sports Medicine Position Stand. Appropriate physical activity intervention strategies for weight loss and prevention of weight regain for adults. *Med Sci Sports Exerc* 2009; 41: 459–71.
15. Jakicic JM. The role of physical activity in prevention and treatment of body weight gain in adults. *J Nutr* 2002; 132: 3826S–9S.
16. Shiraev T, Barclay G. Evidence based exercise – clinical benefits of high intensity interval training. *Aust Fam Physician* 2012; 41: 960–2.

Literatur zu Kapitel 15 Metabolisches Syndrom

1. Reaven GM. Role of insulin resistance in human disease. *Diabetes* 1988; 37: 1595–607.
2. Das UN. Metabolic syndrome X: an inflammatory condition? *Curr Hypertens Rep* 2004; 6: 66–73.
3. Ford ES, Giles WH, Dietz WH. Prevalence of the metabolic syndrome among US adults: findings from the third National Health and Nutrition Examination Survey. *JAMA* 2002; 287: 356–9.
4. Galassi A, Reynolds K, He J. Metabolic syndrome and risk of cardiovascular disease: a meta-analysis. *Am J Med* 2006; 119: 812–9.
5. König D, Bonner G, Berg A. (The role of adiposity and inactivity in primary prevention of cardiovascular disease). *Herz* 2007; 32: 553–9.
6. Lakka HM, Laaksonen DE, Lakka TA, et al. The metabolic syndrome and total and cardiovascular disease mortality in middle-aged men. *JAMA* 2002; 288: 2709–16.
7. Lin CH, Chiang SL, Tzeng WC, Chiang LC. Systematic review of impact of lifestyle-modification programs on metabolic risks and patient-reported outcomes in adults with metabolic syndrome. *Worldviews Evid Based Nurs* 2014; 11: 361–8.
8. Owens S, Galloway R. Childhood obesity and the metabolic syndrome. *Curr Atheroscler Rep* 2014; 16: 436.
9. Ravussin E, Smith SR. Increased fat intake, impaired fat oxidation, and failure of fat cell proliferation result in ectopic fat storage, insulin resistance, and type 2 diabetes mellitus. *Ann N Y Acad Sci* 2002; 967: 363–78.
10. Grundy SM. Pre-diabetes, metabolic syndrome, and cardiovascular risk. *J Am Coll Cardiol* 2012; 59: 635–43.
11. Savage DB, Petersen KF, Shulman GI. Mechanisms of insulin resistance in humans and possible links with inflammation. *Hypertension* 2005; 45: 828–33.
12. Unger RH. Lipotoxic diseases. *Annu Rev Med* 2002; 53: 319–36.
13. Brady MJ. IRS2 takes center stage in the development of type 2 diabetes. *J Clin Invest* 2004; 114: 886–8.
14. Steinberg HO, Paradisi G, Hook G, Crowder K, Cronin J, Baron AD. Free fatty acid elevation impairs insulin-mediated vasodilation and nitric oxide production. *Diabetes* 2000; 49: 1231–8.
15. Al Dokhi LM. Adipokines and etiopathology of metabolic disorders. *Saudi Med J* 2009; 30: 1123–32.
16. Christensen JO, Svendsen OL, Hassager C, Christiansen C. Leptin in overweight postmenopausal women: no relationship with metabolic syndrome X or effect of exercise in addition to diet. *Int J Obes Relat Metab Disord* 1998; 22: 195–9.
17. Wajchenberg BL. Subcutaneous and visceral adipose tissue: their relation to the metabolic syndrome. *Endocr Rev* 2000; 21: 697–738.
18. Beltowski J. Adiponectin and resistin-new hormones of white adipose tissue. *Med Sci Monit* 2003; 9: RA55–RA61.

19. Pedersen BK, Saltin B. Exercise as medicine – evidence for prescribing exercise as therapy in 26 different chronic diseases. *Scand J Med Sci Sports* 2015; 25 Suppl 3: 1–72.
20. Lee S, Kim Y. Effects of exercise alone on insulin sensitivity and glucose tolerance in obese youth. *Diabetes Metab J* 2013; 37: 225–32.
21. Bassi N, Karagodin I, Wang S, et al. Lifestyle modification for metabolic syndrome: a systematic review. *Am J Med* 2014; 127: 1242–10.
22. Andersen CJ, Fernandez ML. Dietary strategies to reduce metabolic syndrome. *Rev Endocr Metab Disord* 2013; 14: 241–54.
23. Gordon B, Chen S, Durstine JL. The effects of exercise training on the traditional lipid profile and beyond. *Curr Sports Med Rep* 2014; 13: 253–9.
24. König D, Deibert P, Dickhuth HH, Berg A. Physical activity and dyslipoproteinemia. *MMW Fortschr Med* 2004; 146: 34–7.
25. König D, Deibert P, Dickhuth HH, Berg A. Bewegungstherapie bei Diabetes mellitus Typ II – metabolische Grundlagen und evidenzbasierte Empfehlungen. *Dtsch Z Sportmed* 2010; 57: 242–7.
26. Hwang CL, Wu YT, Chou CH. Effect of aerobic interval training on exercise capacity and metabolic risk factors in people with cardiometabolic disorders: a meta-analysis. *J Cardiopulm Rehabil Prev* 2011; 31: 378–85.
27. Montesi L, Moscattiello S, Malavolti M, Marzocchi R, Marchesini G. Physical activity for the prevention and treatment of metabolic disorders. *Intern Emerg Med* 2013; 8: 655–66.
28. Pattyn N, Cornelissen VA, Eshghi SR, Vanhees L. The effect of exercise on the cardiovascular risk factors constituting the metabolic syndrome: a meta-analysis of controlled trials. *Sports Med* 2013; 43: 121–33.
29. König D, Berg A. (Physical exercise as treatment of type 2 diabetes mellitus). *Internist (Berl)* 2012; 53: 678–87.
30. Carroll S, Dudfield M. What is the relationship between exercise and metabolic abnormalities? A review of the metabolic syndrome. *Sports Med* 2004; 34: 371–418.

Literatur zu Kapitel 16 Dyslipidämie

1. Scheidt-Nave C, Du Y, Knopf H, Schienkiewitz A, Ziese T et al. Verbreitung von Fettstoffwechselstörungen bei Erwachsenen in Deutschland. Ergebnisse der Studie zur Gesundheit Erwachsener in Deutschland (DEGS1). *Bundesgesundheitsbl* 2013; 56: 661–7.
2. Baghshomali S, Bushnell C. Reducing stroke in women with risk factor management: blood pressure and cholesterol. *Womens Health* 2014; 10: 535–44.
3. Reimers CD, Knapp G (2016) Dyslipidämien. In: Mooren F, Knapp G, Reimers CD (Hrsg.) Prävention und Therapie durch Sport. Band 4. Innere Medizin/Pädiatrie. 2. Aufl., Urban & Fischer, München, S. 29–92.
4. Herdegen T. Lipidsenker. In: Herdegen T (Hrsg.) Kurzlehrbuch Pharmakologie und Toxikologie. 3. Aufl., Stuttgart: Thieme, 2014, S. 263–8.
5. Sudhop T, von Bergmann K. Pharmaka zur Therapie von Stoffwechselkrankheiten. In: Schmidt H (Hrsg.) Pharmakologie und Toxikologie für Studium und Praxis. 6. Aufl., Stuttgart, New York: Schattauer, 2007, S. 759–75.
6. Kodama S, Tanaka S, Saito K, Shu M, Sone Y et al. Effect of aerobic exercise training on serum levels of high-density lipoprotein cholesterol: a meta-analysis. *Arch Intern Med* 2007; 167: 999–1008.
7. García-Hermoso A, Ramírez-Vélez R, Ramírez-Campillo R, Peterson MD, Martínez-Vizcaino V. Concurrent aerobic plus resistance exercise versus aerobic exercise alone to improve health outcomes in paediatric obesity: a systematic review and meta-analysis. *Br J Sports Med* 2016 Dec 16. pii: bjsports-2016-096605. doi: 10.1136/bjsports-2016-096605. (Epub ahead of print).

Literatur zu Kapitel 17 Chronische Nierenerkrankungen

1. Ojo A. Addressing the Global Burden of Chronic Kidney Disease Through Clinical and Translational Research. *Transactions of the American Clinical and Climatological Association* 2014; 125: 229–246.
2. Heiwe S, Jacobson SH. Exercise training for adults with chronic kidney disease. *Cochrane Database Syst Rev* 2011 Oct 5; 10.
3. Beddhu S, Baird BC, Zitterkoph J, Neilson J, Greene T. Physical activity and mortality in chronic kidney disease (NHANES III). *Clin J Am Soc Nephrol* 2009; 4: 1901–1906.
4. Tentori F, Elder SJ, Thumma J et al. Physical exercise among participants in the Dialysis Outcomes and Practice Patterns Study (DOPPS): correlates and associated outcomes. *Nephrol Dial Transplant* 2010; 25: 3050–3062.
5. Painter P. Physical functioning in end-stage renal disease patients: update 2005. *Hemodial Int* 2005; 9: 218–235.
6. Vanholder R, Massy Z, Argiles A, Spasovski G, Verbeke F, Lameire N. Chronic kidney disease as cause of cardiovascular morbidity and mortality. *Nephrol Dial Transplant* 2005; 20: 1048–1056.

7. Finkelstein FO, Wuerth D, Finkelstein SH. Health related quality of life and the CKD patient: challenges for the nephrology community. *Kidney Int* 2009; 76: 946–952.
8. Moinuddin I, Leehey DJ. A comparison of aerobic exercise and resistance training in patients with and without chronic kidney disease. *Adv Chronic Kidney Dis* 2008; 15: 83–96.
9. Johansen KL, Painter P. Exercise in individuals with CKD. *Am J Kidney Dis*. 2012 Jan; 59: 126–34.
10. Daul AE. Körperliches Training und Dialyse. *Der Nephrologe* 2011, 6: 537–547.
11. Heiwe S, Jacobson SH. Exercise training in adults with CKD: a systematic review and meta-analysis. *Am J Kidney Dis* 2014 Sep; 64: 383–937.
12. Smart N, Steele M. Exercise training in haemodialysis patients: a systematic review and meta-analysis. *Nephrology (Carlton)*. 2011 Sep; 16: 626–32.
13. Smart N, McFarlane J, Cornelissen V. The Effect of Exercise Therapy on Physical Function, Biochemistry and Dialysis Adequacy in Haemodialysis Patients: A Systematic Review and Meta-Analysis. *Open Journal of Nephrology* 2013, 3: 25–36.
14. Vaithilingam I, Polkinghorne KR, Atkins RC, Kerr PG. Time and exercise improve phosphate removal in hemodialysis patients. *Am J Kidney Dis* 2004; 43: 85–89.
15. Anding K, Fuhrmann I. Praktische Umsetzung von Sporttherapie während der Hämodialyse mit dem Sächsischen Strukturierten Sporttherapie Programm (SSSP). *Nieren- und Hochdruckkrankheiten* 2015; 44: 375–382.
16. Anding K, Bär T, Trojniak-Hennig J, et al. A structured exercise programme during haemodialysis for patients with chronic kidney disease: clinical benefit and long-term adherence. *BMJ Open* 2015; 5: e008709. doi:10.1136/bmjopen-2015-008709.

Literatur zu Kapitel 18 Lebererkrankungen

1. de Boer A1, Kluft C, Kroon JM, Kasper FJ, Schoemaker HC et al. Liver blood flow as a major determinant of the clearance of recombinant human tissue-type plasminogen activator. *Thromb Haemost* 1992; 67: 83–7.
2. Suh SH, Paik IY, Jacobs K. Regulation of blood glucose homeostasis during prolonged exercise. *Mol Cells* 2007; 23: 272–9.
3. McKenna O, Cunningham C, Gissane C, Blake C. Management of the extrahepatic symptoms of chronic hepatitis C. *Am J Phys Med Rehabil* 2013; 92: 504–12.
4. Caldwell S, Lazo M. Is exercise an effective treatment for NASH? Knowns and unknowns. *Ann Hepatol* 2009; 8: 60–6.
5. Harrington D. Viral hepatitis and exercise. *Med. Sci. Sports Exerc* 2000; 32: 422–30.
6. Chalmers T, Eckhardt R, Reynolds W, Cigarroa J, Deane N et al. The treatment of acute infectious hepatitis. *Journal of Clinical Investigation* 1955; 34: 1163–1235.
7. Repsher L, Freeborn R. Effects of early and vigorous exercise on recovery from infectious hepatitis. *New England Journal of Medicine* 1969; 281: 1393–96.
8. Wiese M, Metzner C, Ziesenhenn K, Haupt R. Effect of a bicycle ergometry training program in the acute phase of viral hepatitis on its late prognosis. *Zeitschrift für Gesamte Innere Medizin* 1983; 38: 42–44.
9. Graubaus H, Metzner C, Ziesenhenn K. Körperliche Belastung und Hepatitisverlauf. *Deutsche Medizinische Wochenschrift* 1987; 112: 47–49.
10. Ishida A, Sumiya N, Ueno F. The effects of physical activity on rehabilitation for acute hepatitis. *Tokai Journal of Experimental and Clinical Medicine* 1996; 21: 1–6.
11. Ritland S, Petlund C, Knudsen T, Skrede S. Improvement of physical capacity after long-term training in patients with chronic active hepatitis. *Scandinavian Journal of Gastroenterology* 1983; 18: 1083–7.
12. Moon J1, Kallman J, Winter P, Srishord M, Fang Y et al. Disparities in activity level and nutrition between patients with chronic hepatitis C and blood donors. *PM R* 2012; 4: 436–41.
13. Vázquez-Vandyck M, Roman S, Vázquez JL, Huacuja L, Khalsa G et al. Effect of Breathwalk on body composition, metabolic and mood state in chronic hepatitis C patients with insulin resistance syndrome. *World J Gastroenterol* 2007; 13: 6213–8.
14. Payen J, Pillard F, Mascarell V, Riviere D, Couzigou P et al. Is physical activity possible and beneficial for patients with hepatitis C receiving pegylated interferon and ribavarin therapy? *Gastroenterologie Clinique et Biologique* 2009; 33: 8–14.
15. Konishi I, Hiasa Y, Tokumoto Y, Abe M, Furukawa S et al. Aerobic exercise improves insulin resistance and decreases body fat and serum levels of leptin in patients with hepatitis C virus. *Hepatology Research* 2011; 41: 928–35.
16. Pattullo V, Duarte-Rojo A, Soliman W, Vargas-Vorackova F, Sockalingam S, et al. A 24-week dietary and physical activity lifestyle intervention reduces hepatic insulin resistance in the obese with chronic hepatitis C. *Liver International* 2012; 33: 410–9.

17. Abd El-Kader SM, Al-Jiffri OH, Al-Shreef FM. Liver enzymes and psychological well-being response to aerobic exercise training in patients with chronic hepatitis C. *Afr Health Sci* 2014; 14: 414–9.
18. Church TS, Kuk JL, Ross R, Priest EL, Biloft E et al. Association of cardiorespiratory fitness, body mass index, and waist circumference to nonalcoholic fatty liver disease. *Gastroenterology* 2006; 130: 2023–30.
19. Goncalves I, Oliveira P, Ascensao A, Magalhaes J. Exercise as a therapeutic tool to prevent mitochondrial degeneration in nonalcoholic steatohepatitis. *Europ J of Clinical Investigation* 2013; 43: 1184–94.
20. McMillan KP1, Kuk JL, Church TS, Blair SN, Ross R. Independent associations between liver fat, visceral adipose tissue, and metabolic risk factors in men. *Appl Physiol Nutr Metab* 2007; 32: 265–72.
21. Krasnoff JB, Painter PL, Wallace JP, Bass NM, Merriman RB. Health-related fitness and physical activity in patients with nonalcoholic fatty liver disease. *Hepatology* 2008; 47: 1158–66.
22. Johnson N, Sachinwalla T, Walton D, Smith K, Armstrong A et al. Aerobic exercise training reduces hepatic and visceral lipids in obese individuals without weight loss. *Journal of Hepatology* 2009; 50: 1105–12.
23. Alameri H, Sanai F, Dukhayil M, Azzam N, Al-Swat K et al. A. Six minute walk test to assess functional capacity in chronic liver disease patients. *World Journal of Gastroenterology* 2007; 13: 3996–4001.
24. Galant L, Forgiarini L, Dias A, Marroni C. Functional status, respiratory muscle strength and quality of life in patients with cirrhosis. *Brazilian Journal of Physical Therapy* 2012; 16: 30–4.
25. Terziyski K, Andonov V, Marinov B, Konstianev S. Exercise performance and ventilatory efficiency in patients with mild and moderate liver cirrhosis. *Clinical and Experimental Pharmacology and Physiology* 2008; 35: 135–40.
26. Wiesinger GF, Quittan M, Zimmermann K, Nuhr M, Wichlas M et al. Physical performance and health-related quality of life in men on a liver transplantation waiting list. *J Rehabil Med* 2001; 33: 260–5.
27. Panzak G1, Tarter R, Murali S, Switala J, Lu S et al. Isometric muscle strength in alcoholic and nonalcoholic liver-transplantation candidates. *Am J Drug Alcohol Abuse* 1998; 24: 499–512.
28. Andersen H, Borre M, Jakobsen J, Andersen PH, Vilstrup H. Decreased muscle strength in patients with alcoholic liver cirrhosis in relation to nutritional status, alcohol abstinence, liver function, and neuropathy. *Hepatology* 1998; 27: 1200–6.
29. Casaburi R, Oi S. Effect of liver disease on the kinetics of lactate removal after heavy exercise. *Eur J Appl Physiol* 1989; 59: 89–97.

Literatur zu Kapitel 19 Rheumatoide Arthritis

1. Fenton SA, Kitas GD. Rheumatoid arthritis: Sedentary behaviour in RA – a new research agenda. *Nat Rev Rheumatol* 2016 Dec; 12: 698–700.
2. Krüger K, Mooren FC. Sport, Immunsystem und rheumatologische Erkrankungen. *Akt Rheumatol* 2012; 37: 189–195.
3. Lefèvre S, Knedla A, Tennie C, Kampmann A, Wunrau C, Dinser R et al. Synovial fibroblasts spread rheumatoid arthritis to unaffected joints. *Nat Med* 2009 Dec; 15: 1414–20.
4. Siu S, Haraoui B, Bissonnette R, Bessette L, Roubille C, Richer V, Starnino T, McCourt C, McFarlane A, Fleming P, Kraft J, Lynde C, Gulliver W, Keeling S, Dutz J, Pope JE. A meta-analysis of tumor necrosis factor inhibitors and glucocorticoids on bone density in rheumatoid arthritis and ankylosing spondylitis trials. *Arthritis Care Res (Hoboken)*. 2014 Nov 21.
5. Lemmey AB, Marcora SM, Chester K, Wilson S, Casanova F, Maddison PJ. Effects of high-intensity resistance training in patients with rheumatoid arthritis: a randomized controlled trial. *Arthritis Rheum* 2009 Dec 15; 61: 1726–34.
6. de Jong Z, Munneke M, Jansen LM, Ronda K, van Schaardenburg DJ, Brand R et al. Differences between participants and nonparticipants in an exercise trial for adults with rheumatoid arthritis. *Arthritis Rheum* 2004 Aug 15; 51: 593–600.
7. Neuberger GB, Aaronson LS, Gajewski B, Embretson SE, Cagle PE, Loudon JK, Miller PA. Predictors of exercise and effects of exercise on symptoms, function, aerobic fitness, and disease outcomes of rheumatoid arthritis. *Arthritis Rheum* 2007 Aug 15; 57: 943–52.
8. Baillet A, Vaillant M, Guinot M, Juvin R, Gaudin P. Efficacy of resistance exercises in rheumatoid arthritis: meta-analysis of randomized controlled trials. *Rheumatology* 2012 Mar; 51: 519–27.
9. Baillet A, Zeboulon N, Gossec L, Combesse C, Bodin LA, Juvin R, Dougados M, Gaudin P. Efficacy of cardiorespiratory aerobic exercise in rheumatoid arthritis: meta-analysis of randomized controlled trials. *Arthritis Care Res* 2010 Jul; 62: 984–92.
10. Pedersen BK, Saltin B. Exercise as medicine – evidence for prescribing exercise as therapy in 26 different chronic diseases. *Scand J Med Sci Sports* 2015 Dec; 25 Suppl 3: 1–72.
11. Shin JH, Lee Y, Kim SG, Choi BY, Lee HS, Bang SY. The beneficial effects of Tai Chi exercise on endothelial function and arterial stiffness in elderly women with rheumatoid arthritis. *Arthritis Res Ther* 2015 Dec 24; 17: 380.
12. Wadley AJ, Veldhuijzen van Zanten JJ, Stavropoulos-Kalinoglou A, Metsios GS, Smith JP, Kitas GD, Aldred S. Three months of moderate-intensity exercise reduced plasma 3-nitrotyrosine in rheumatoid arthritis patients. *Eur J Appl Physiol* 2014 Jul; 114: 1483–92.

13. Brosseau L, Wells GA and Ottawa Panel. Evidence-based clinical practice guidelines for therapeutic exercises in the management of rheumatoid arthritis in adults. *Phys Ther* 2004 Oct; 84: 934–72.
14. Metsios GS, Stavropoulos-Kalinoglou A, Panoulas VF, Wilson M, Nevill AM, Koutedakis Y, Kitas GD. Association of physical inactivity with increased cardiovascular risk in patients with rheumatoid arthritis. *Eur J Cardiovasc Prev Rehabil* 2009 Apr; 16: 188–94.

Literatur zu Kapitel 20 Spondylitis ankylosans

1. Raychaudhuri SP, Deodhar A. The classification and diagnostic criteria of ankylosing spondylitis. *J Autoimmun* 2014 Feb–Mar; 48–49: 128–33.
2. Tsui FW, Tsui HW, Akram A, Haroon N, Inman RD. The genetic basis of ankylosing spondylitis: new insights into disease pathogenesis. *Appl Clin Genet* 2014 May 22; 7: 105–15.
3. Bowness P. HLA B27 in health and disease: a double-edged sword? *Rheumatology* 2002 Aug; 41: 857–68.
4. Smith JA. Update on ankylosing spondylitis: current concepts in pathogenesis. *Curr Allergy Asthma Rep* 2015 Jan; 15: 489.
5. Lubrano E, Spadaro A, Amato G, Benucci M, Cavazzana I, Chimenti MS et al. Tumour necrosis factor alpha inhibitor therapy and rehabilitation for the treatment of ankylosing spondylitis: A systematic review. *Semin Arthritis Rheum* 2015 Apr; 44 :542–550.
6. Millner JR, Barron JS, Beinke KM, Butterworth RH, Chasle BE, Dutton LJ et al. Exercise for ankylosing spondylitis: An evidence-based consensus statement. *Semin Arthritis Rheum* 2016 Feb; 45: 411–27.
7. Aytekin E, Caglar NS, Ozgonenel L, Tutun S, Demiryontar DY, Demir SE. Home-based exercise therapy in patients with ankylosing spondylitis: effects on pain, mobility, disease activity, quality of life, and respiratory functions. *Clin Rheumatol* 2012 Jan; 31: 91–7.
8. O'Dwyer T, O'Shea F, Wilson F. Exercise therapy for spondyloarthritis: a systematic review. *Rheumatol Int* 2014 Jul; 34: 887–902.

Literatur zu Kapitel 21 Infekte des oberen Respirationstrakts

1. Mäkelä MJ, Puhakka T, Ruuskanen O, Leinonen M, Saikku P, Kimpimäki M et al. Viruses and bacteria in the etiology of the common cold. *J Clin Microbiol* 1998; 36: 539–42.
2. Monto AS. Studies of the community and family: acute respiratory illness and infection. *Epidemiol Rev* 1994; 16: 351–73.
3. Nieman DC. Exercise, Infection, and Immunity. *Int J Sports Med* 1994; 15: S131–S141.
4. Nieman DC, Johanssen LM, Lee JW, Arabatzis K. Infectious episodes in runners before and after the Los Angeles Marathon. *J Sports Med Phys Fitness* 1990; 30: 316–28.
5. Nieman DC, Henson DA, Dumke CL, Lind RH, Shooter LR, Gross SJ. Relationship between salivary IgA secretion and upper respiratory tract infection following a 160-km race. *J Sports Med Phys Fitness* 2006; 46: 158–62.
6. Gleeson M, McDonald WA, Pyne DB, Cripps AW, Francis JL, Fricker PA et al. Salivary IgA levels and infection risk in elite swimmers. *Med Sci Sports Exerc* 1999; 31: 67–73.
7. Gleeson M, Bishop N, Oliveira M, Tauler P. Influence of training load on upper respiratory tract infection incidence and antigen-stimulated cytokine production. *Scand J Med Sci Sports* 2013; 23: 451–7.
8. Neville V, Gleeson M, Folland JP. Salivary IgA as a risk factor for upper respiratory infections in elite professional athletes. *Med Sci Sports Exerc* 2008; 40: 1228–36.
9. Novas AMP, Rowbottom DG, Jenkins DG. Tennis, incidence of URTI and salivary IgA. *Int J Sports Med* 2003; 24: 223–9.
10. Robson-Ansley P, Howatson G, Tallent J, Mitcheson K, Walshe I, Toms C et al. Prevalence of allergy and upper respiratory tract symptoms in runners of the London marathon. *Med Sci Sports Exerc* 2012; 44: 999–1004.
11. Pyne DB, Baker MS, Fricker PA, McDonald WA, Telford RD, Weidemann MJ. Effects of an intensive 12- wk training program by elite swimmers on neutrophil oxidative activity. *Med Sci Sports Exerc* 1995; 27: 536–42.
12. Pyne DB, McDonald WA, Gleeson M, Flanagan A, Clancy RL, Fricker PA. Mucosal immunity, respiratory illness, and competitive performance in elite swimmers. *Med Sci Sports Exerc* 2001; 33: 348–53.
13. Nieman DC, Johanssen LM, Lee JW. Infectious episodes in runners before and after a roadrace. *J Sports Med Phys Fitness* 1989; 29: 289–96.
14. Matthews CE, Ockene IS, Freedson PS, Rosal MC, Merriam PA, Herbert JR. Moderate to vigorous physical activity and risk of upper-respiratory tract infection. *Med Sci Sports Exerc* 2002; 34: 1242–8.
15. Kekkonen RA, Vasankari TJ, Vuorimaa T, Haahtela T, Julkunen I, Korpela R. The effect of probiotics on respiratory infections and gastrointestinal symptoms during training in marathon runners. *Int J Sport Nutr Exerc Metab* 2007; 17: 352–63.

16. Henson D, Nieman D, Davis JM, Dumke C, Gross S, Murphy A et al. Post-160-km race illness rates and decreases in granulocyte respiratory burst and salivary IgA output are not countered by quercetin ingestion. *Int J Sports Med* 2008; 29: 856–63.
17. Peters EM, Goetzsche JM, Grobbelaar B, Noakes TD. Vitamin C supplementation reduces the incidence of posttrace symptoms of upper-respiratory-tract infection in ultramarathon runners. *Am J Clin Nutr* 1993; 57: 170–4.
18. Himmelstein SA, Robergs RA, Koehler KM, Lewis SL, Qualls CR. Vitamin C supplementation and upper respiratory tract infections in marathon runners. *J Exerc Physiol* 1998; 1: 1–21.
19. Nieman DC, Peters EM, Henson DA, Nevines EI, Thompson MM. Influence of vitamin C supplementation on cytokine changes following an ultramarathon. *J Interferon Cytokine Res* 2000; 20: 1029–35.
20. Walsh NP, Gleeson M, Pyne DB, Nieman DC, Dhabhar FS, Shephard RJ et al. Position statement. Part two: Maintaining immune health. *Exerc Immunol Rev* 2011; 17: 64–103.

Literatur zu Kapitel 22 HIV-Infektion

1. Robert-Koch-Institut. Zum Welt-AIDS-Tag. Schätzung der Prävalenz und Inzidenz von HIV-Infektionen in Deutschland, Stand Ende 2012. *Epidemiologisches Bulletin* 2012; 47: 465–480.
2. Cavarelli M, Scarlatti G. HIV-1 infection: the role of the gastrointestinal tract. *Am J Reprod Immunol* 2014; 71: 537–542.
3. Raso V, Shephard RJ, Casseb JS, Duarte AJ, Greve JM. Aerobic power and muscle strength of individuals living with HIV/AIDS. *J Sports Med Phys Fitness* 2014; 54: 100–107.
4. Engelson ES, Agin D, Kenya S, Werber-Zion G, Luty B, Albu et al. Body composition and metabolic effects of a diet and exercise weight loss regimen on obese, HIV-infected women. *Metabolism* 2006; 55: 1327–1336.
5. MacArthur RD, Levine SD, Birk TJ. Supervised exercise training improves cardiopulmonary fitness in HIV-infected persons. *Med Sci Sports Exerc* 1993; 25: 684–688.
6. Mutimura E, Crowther NJ, Cade TW, Yarasheski KE, Stewart A. Exercise training reduces central adiposity and improves metabolic indices in HAART-treated HIV-positive subjects in Rwanda: a randomized controlled trial. *AIDS Res Hum Retroviruses* 2008; 24: 15–23.
7. LaPerriere A, Antoni MH, Schneiderman N, Ironson G, Klimas N et al. Exercise intervention attenuates emotional distress and natural killer cell decrements following notification of positive serologic status for HIV-1. *Biofeedback Self Regul* 1990; 15: 229–242.
8. Ahmad B, Glufke K, Grau M, Sandig D, Rockstroh J, Vogel M et al. Influence of endurance training and marathon running on red cell deformability in HIV patients. *Clin Hemorheol Microcirc* 2013; DOI 10.3233.
9. Perna FM, LaPerriere A, Klimas N, Ironson G, Perry A, Pavone J et al. Cardiopulmonary and CD4 cell changes in response to exercise training in early symptomatic HIV infection. *Med. Sci. Sports Exerc* 1999; 31: 973–979.
10. Stringer WW, Berezovskaya M, O'Brien WA, Beck CK, Casaburi R. The effect of exercise training on aerobic fitness, immune indices, and quality of life in HIV+ patients. *Med Sci Sports Exerc* 1998; 30: 11–16.
11. Terry L, Sprinz E, Ribeiro JP. Moderate and high intensity exercise training in HIV-1 seropositive individuals: a randomized trial. *Int J Sports Med* 1999; 20: 142–146.
12. Schlenzig CA. Sport und HIV-Infektion. Reaktionen des Immunsystems und der Psyche HIV-Infizierter und AIDS-Patienten auf kontrollierte sporttherapeutische Langzeitbehandlungen. Dissertation, Universität Heidelberg 1992.
13. Smith BA, Neidig JL, Nickel JT, Mitchell GL, Para MF, Fass RJ. Aerobic exercise: effects on parameters related to fatigue, dyspnea, weight and body composition in HIV-infected adults. *AIDS* 2001; 15: 693–701.
14. Vaiyapuri A, Ivor P, Jagatheesan A, Durai P, Rajendran K. Effect of aerobic exercise training on cardio vascular fitness, quality of life and CD4 count in people with HIV/AIDS. *Int J Adv Lif Sc* 2013; 6: 569–574.
15. Antczak AW. Effekte eines aeroben Ausdauertrainings auf HIV-positive Patienten unter antiretroviraler Therapie. Dissertation, Freie Universität Berlin, Medizinische Fakultät Charité 2012.
16. LaPerriere A, Antoni MH, Ironson G, Perry A, McCabe P, Klimas N. Effects of aerobic exercise training on lymphocyte subpopulations. *Int J Sports Med* 1994; 15: 127–130.
17. Lox CL, McAuley E, Tucker RS. Aerobic and resistance exercise training effects on body composition, muscular strength, and cardiovascular fitness in an HIV-1 population. *Int J Behav Med* 1996; 3: 55–69.
18. Thöni GJ, Fedou C, Brun JF, Fabre J, Renard E, Reynes J et al. Reduction of fat accumulation and lipid disorders by individualized light aerobic training in human immunodeficiency virus infected patients with lipodystrophy and/or dyslipidemia. *Diabetes Metab* 2002; 28: 397–404.
19. Terry L, Sprinz E, Stein R, Medeiros NB, Oliveira J, Ribeiro JP. Exercise training in HIV-1-infected individuals with dyslipidemia and lipodystrophy. *Med Sci Sports Exerc* 2006; 38: 411–417.

20. Lindegaard B, Hansen T, Hvid T, van Hal, G, Plomgaard P, Ditlevsen S et al. The effect of strength and endurance training on insulin sensitivity and fat distribution in human immunodeficiency virus-infected patients with lipodystrophy. *J Clin Endocrinol Metab* 2008; 93: 3860–3869.
21. Roubenoff R, Weiss L, McDermott A, Heflin T, Cloutier G J, Wood M, Gorbach S. A pilot study of exercise training to reduce trunk fat in adults with HIV-associated fat redistribution. *AIDS* 1999; 13: 1373–1375.
22. Spence DW, Galantin M L, Mossberg K A, Zimmerman S O. Progressive resistance exercise: effect on muscle function and anthropometry of a select AIDS population. *Arch Phys Med Rehabil* 1990; 71: 644–648.
23. de Souza P M L, Jacob-Filho W, Santarem J M, Zomignan A A, Burattini M N. Effect of progressive resistance exercise on strength evolution of elderly patients living with HIV compared to healthy controls. *Clinics* 2010; 66: 261–266.
24. Yarasheski K E, Tebas P, Stanerson E, Claxton S, Marin D, Bae K et al. Resistance exercise training reduces hypertriglyceridemia in HIV-infected men treated with antiviral therapy. *J Appl Physiol* 2001; 90: 133–138.
25. Roubenoff R, McDermott A, Weiss L, Suri J, Wood M, Bloch R, Gorbach S. Short-term progressive resistance training increases strength and lean body mass in adults infected with human immunodeficiency virus. *AIDS* 1999; 13: 231–239.
26. Robinson F P, Quinn L T, Rimmer J H. Effects of high-intensity endurance and resistance exercise on HIV metabolic abnormalities: a pilot study. *Biol Res Nurs* 2007; 8: 177–185.
27. Rojas Salcedo R I. Vergleich zwischen den Auswirkungen eines Sport- und eines psychotherapeutischen Programms auf die Lebensqualität HIV-positiver Menschen. Dissertation. Regensburg: S. Roderer, 2001.
28. Hand G A, Phillips K D, Dudgeon W D, William Lyerly G, Larry Durstine J, Burgess S E. Moderate intensity exercise training reverses functional aerobic impairment in HIV-infected individuals. *AIDS Care* 2008; 20: 1066–1074.
29. Fillipas S, Oldmeadow L B, Bailey M J, Cherry C L. A six-month, supervised, aerobic and resistance exercise program improves self-efficacy in people with human immunodeficiency virus: a randomised controlled trial. *Aust J Physiother* 2006; 52: 185–190.
30. Dolan S E, Frontera W, Librizzi J, Ljungquist K, Juan S, Dorman R et al. Effects of a supervised home-based aerobic and progressive resistance training regimen in women infected with human immunodeficiency virus: a randomized trial. *Arch Intern Med* 2006; 166: 1225–1231.
31. Rigsby L W, Dishman R K, Jackson A W, Maclean G S, Raven P B. Effects of exercise training on men seropositive for the human immunodeficiency virus-1. *Med. Sci. Sports Exerc* 1992; 24: 6–12.
32. Farinatti P T, Borges J P, Gomes R D, Lima D, Fleck S J. Effects of a supervised exercise program on the physical fitness and immunological function of HIV-infected patients. *J Sports Med Phys Fitness* 2010; 50: 511–518.
33. Baigis J, Korniewicz D M, Chase G, Butz A, Jacobson D, Wu A W. Effectiveness of a home-based exercise intervention for HIV-infected adults: a randomized trial. *J Assoc Nurses AIDS Care* 2002; 13: 33–45.
34. Borges J, Soares P, Farinatti P. Autonomic modulation following exercise is impaired in HIV patients. *Int J Sports Med* 2012; 33: 320–24.
35. Deresz L F, Sprinz E, Kramer A S, Cunha G, De Oliveira A R, Sporleder H et al. Regulation of oxidative stress in response to acute aerobic and resistance exercise in HIV-infected subjects: A case-control study. *AIDS Care – Psychological and Socio-Medical Aspects of AIDS/HIV* 2010; 22: 1410–1417.
36. Roubenoff R, Wilson J B. Effect of resistance training on self-reported physical functioning in HIV infection. *Med Sci Sports Exerc* 2001; 33: 1811–17.
37. Galantino M L, Shepard K, Krafft L, LaPerriere A, Ducette J, Sorbello A et al. The effect of group aerobic exercise and t'ai chi on functional outcomes and quality of life for persons living with acquired immunodeficiency syndrome. *Journal of Alternative and Complementary Medicine* 2005; 11: 1085–92.
38. LaPerriere A, Klimas N, Major P, Perry A. Acquired Immune Deficiency Syndrome. In: Durstine J, Bloomquist L, Painter S, Pope C (Hrsg.): *Exercise Management for Persons with Chronic Disease and Disabilities*. American college of sports medicine, human kinetics 1998, S. 132–36.
39. Mooren F, Fink K (2016) Sport und HIV-Erkrankung. In Mooren F, Knapp, Reimers CD (Hrsg.) *Prävention und Therapie durch Sport*. München: Urban & Fischer, Band 3, S. 205–50.

Literatur zu Kapitel 23 Asthma bronchiale

1. Global strategy for asthma management and prevention. [cited 2016.]
2. Sears M R. Trends in the prevalence of asthma. *Chest* 2014; 145: 219–25.
3. Reddel, H., et al., Differences between asthma exacerbations and poor asthma control. *Lancet* 1999; 353: 364–9.
4. Masoli M et al. The global burden of asthma: executive summary of the GINA Dissemination Committee report. *Allergy* 2004; 59: 469–78.
5. Beasley R, Semprin A, Mitchell E. A. Risk factors for asthma: is prevention possible? *Lancet* 2015; 386: 1075–85.
6. Busse W. W, Lemanske Jr. R F. Asthma. *N Engl J Med* 2001; 344: 350–62.

7. Anderson SD, Kippelen P. Assessment and prevention of exercise-induced bronchoconstriction. *Br J Sports Med* 2012; 46: 391–6.
8. Ansley L, Rae G, Hull JH. Practical approach to exercise-induced bronchoconstriction in athletes. *Prim Care Respir J* 2013; 22: 122–5.
9. Parsons JP et al. An official American Thoracic Society clinical practice guideline: exercise-induced bronchoconstriction. *Am J Respir Crit Care Med* 2013; 187: 1016–27.
10. Nationale Versorgungsleitlinie Asthma. [cited Juli 2011; Version 1.3:]
11. Bateman ED, Hurd SS, Barnes PJ, Bousquet J, Drazen JM, FitzGerald M et al. Global strategy for asthma management and prevention. *European Respiratory Journal* 2008; 31: 143–178.
12. Kerstjens HA et al. Influence of treatment on peak expiratory flow and its relation to airway hyperresponsiveness and symptoms. The Dutch CNSLD Study Group. *Thorax* 1994; 49: 1109–15.
13. Anderson SD. Indirect challenge tests: Airway hyperresponsiveness in asthma: its measurement and clinical significance. *Chest* 2010; 138 (2 Suppl): 25S–30S.
14. Bacon SL et al. The effects of a multisite aerobic exercise intervention on asthma morbidity in sedentary adults with asthma: the Ex-asthma study randomised controlled trial protocol. *BMJ Open* 2013; 3: 6.
15. Carson KV et al. Physical training for asthma. *Cochrane Database Syst Rev* 2013; 9: CD001116.
16. Mediano O et al. Dynamic hyperinflation in patients with asthma and exercise-induced bronchoconstriction. *Ann Allergy Asthma Immunol* 2017.
17. Chapman KR et al. Suboptimal asthma control: prevalence, detection and consequences in general practice. *Eur Respir J* 2008; 31: 320–5.
18. Banzer W. Körperliche Aktivität und Gesundheit. Berlin/Heidelberg: Springer 2017.
19. Klijn P et al. Nonlinear exercise training in advanced chronic obstructive pulmonary disease is superior to traditional exercise training. A randomized trial. *Am J Respir Crit Care Med* 2013; 188: 193–200.
20. Freitas PD et al. The Role of Exercise in a Weight-Loss Program on Clinical Control in Obese Adults with Asthma. A Randomized Controlled Trial. *Am J Respir Crit Care Med* 2017; 195: 32–42.
21. Weiner P et al. Inspiratory muscle training in patients with bronchial asthma. *Chest* 1992; 102: 1357–61.

Literatur zu Kapitel 24 Chronische obstruktive Lungenerkrankung

1. GOLD. Available from: <http://www.copdgold.org/>.
2. Barnes PJ. Therapy of chronic obstructive pulmonary disease. *Pharmacol Ther* 2003; 97: 87–94.
3. Maltais F et al. An official American Thoracic Society/European Respiratory Society statement: update on limb muscle dysfunction in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2014; 189: e15–62.
4. Barnes PJ, Celli BR. Systemic manifestations and comorbidities of COPD. *Eur Respir J* 2009; 33: 1165–85.
5. Papi A et al. Infections and airway inflammation in chronic obstructive pulmonary disease severe exacerbations. *Am J Respir Crit Care Med* 2006; 173: 1114–21.
6. Voshaar T et al. COPD: Einsatz inhalativer Mono- und Kombinationstherapien. *Dtsch Arztebl International* 2016; 113: 4.
7. Spruit MA et al. An official American Thoracic Society/European Respiratory Society statement: key concepts and advances in pulmonary rehabilitation. *Am J Respir Crit Care Med* 2013; 188: e13–64.
8. Garosch I, Gloeckl R, Damm E, Schwedhelm AL, Buhrow D, Jerrentrup A, Spruit MA, Kenn K. Short-term effects of supplemental oxygen on 6-minute walk test outcomes in COPD patients – a randomized, placebo-controlled, single-blind, cross-over trial. *Chest* 2016; 8. pii: S0012–3692: p. 62572–2.
9. Levine S et al. COPD elicits remodeling of the diaphragm and vastus lateralis muscles in humans. *J Appl Physiol* (1985), 2013; 114: 1235–45.
10. Gloeckl R, Marinov B, Pitta F. Practical recommendations for exercise training in patients with COPD. *Eur Respir Rev* 2013; 22: 178–86.
11. Lotters F et al. Effects of controlled inspiratory muscle training in patients with COPD: a meta-analysis. *Eur Respir J* 2002; 20: 570–6.
12. Pleguezuelos E et al. Effects of whole body vibration training in patients with severe chronic obstructive pulmonary disease. *Respirology* 2013; 18: 1028–34.
13. Greulich T et al. Benefits of whole body vibration training in patients hospitalised for COPD exacerbations – a randomized clinical trial. *BMC Pulm Med* 2014; 14: 60.
14. Gloeckl R et al. Effects of whole body vibration in patients with chronic obstructive pulmonary disease – a randomized controlled trial. *Respir Med* 2012; 106: 75–83.
15. Sillen MJ et al. Efficacy of lower-limb muscle training modalities in severely dyspnoeic individuals with COPD and quadriceps muscle weakness: results from the DICES trial. *Thorax* 2014; 69: 525–31.

16. Casaburi R, Maltais F, Porszasz J, Albers F, Deng Q, Iqbal A, Paden HA, O'Donnell DE; 205.440 Investigators. Effects of tiotropium on hyperinflation and treadmill exercise tolerance in mild to moderate chronic obstructive pulmonary disease. *Ann Am Thorac Soc* 2014; 11: 1351–61.
17. Segers J et al. Feasibility of neuromuscular electrical stimulation in critically ill patients. *J Crit Care* 2014; 29: 1082–8.
18. Troosters T et al. Resistance training prevents deterioration in quadriceps muscle function during acute exacerbations of chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2010; 181: 1072–7.
19. Handschin C, Spiegelman BM. The role of exercise and PGC1alpha in inflammation and chronic disease. *Nature* 2008; 454: 463–9.
20. Lowie E et al. Clusters of Comorbidities Based on Validated Objective Measurements and Systemic Inflammation in Patients with Chronic Obstructive Pulmonary Disease. *Am J Respir Crit Care Med* 2013; 187: 728–35.
21. Partridge MR, Karlsson N, Small IR. Patient insight into the impact of chronic obstructive pulmonary disease in the morning: an internet survey. *Curr Med Res Opin* 2009; 25: 2043–8.

Literatur zu Kapitel 25 Tumorerkrankungen

1. Robert-Koch-Institut und Gesellschaft der epidemiologischen Krebsregister e. V. (GEKID). Krebs in Deutschland 2011/2012 – Häufigkeiten und Trends 2015, Berlin: 2015.
2. Shah C, Vicini FA. Breast cancer-related arm lymphedema: incidence rates, diagnostic techniques, optimal management and risk reduction strategies. *Int J Radiat Oncol Biol Phys* 2011; 81: 907–14.
3. Pesendorfer LM, Zimmer P, Galvao DA, Zopf EM, Bloch W, Baumann FT. Impact of physical inactivity on the multifactorial process of developing cancer-related cognitive impairment. *Journal of Cancer Science and Therapy* 2016; DOI: 10.4172/1948–5956.1000417.
4. Friedenreich CM. Physical activity and breast cancer: Review of the epidemiologic evidence and biologic mechanisms. *Recent Results Cancer Res* 2011; 188: 25–39.
5. Schmidt ME, Steindorf K, Mutschelknauss E, Slinger T, Obi N et al. Physical activity and postmenopausal breast cancer: Effective periods in life and effect modification by different breast cancer characteristics. *Cancer Epidemiol Biomarkers Prev* 2008; 173: 402–10.
6. Irwin ML, Cartmel B, Gross CP, et al: Randomized exercise trial of aromatase inhibitor induced arthralgia in breast cancer survivors. *J Clin Oncol* 2015; 33: 1104–1111.
7. Mustian K, Peppone L, Darling T, Palesh O, Heckler C, Morrow G. A 4-Week Home-Based Aerobic and Resistance Exercise Program During Radiation Therapy: A Pilot Randomized Clinical Trial. *J Support Oncol* 2009; 7: 158–67.
8. Daley AJ, Crank H, Saxton JM, Mutrie N, Coleman R, Roalfe A. Randomized trial of exercise therapy in women treated for breast cancer. *J Clin Oncol* 2007; 25, 1: 713–21.
9. Culos-Reed SN, Carlson LE, Daroux LM, Hatley-Aldous S. A pilot study of yoga for breast cancer survivors: physical and psychological benefits. *Psychooncology* 2006; 15: 891–7.
10. Schmitz KH, Ahmed RL, Troxel A, Cheville A, Smith R et al. Weight lifting in women with breast-cancer-related lymphedema. *N Engl J Med* 2009; 361: 664–73.
11. Winters-Stone KM, Dobek J, Nail LM, Bennett JA, Leo MC et al. Impact + resistance training improves bone health and body composition in prematurely menopausal breast cancer survivors: a randomized controlled trial. *Osteoporose Int* 2013; 24: 637–46.
12. McKenzie D, Kalda A: Effect of upper extremity exercise on secondary lymphedema in breast cancer patients: a pilot study. *J Clin Oncol* 2003; 21: 463–6.
13. Payne JK, Held J, Thorpe J, Shaw H. Effect of exercise on biomarkers, fatigue, sleep disturbances, and depressive symptoms in older women with breast cancer receiving hormonal therapy. *Oncol Nurs Forum* 2008; 35: 635–42.
14. Fernandez-Lao CC, Cantarero-Villanueva I, Ariza-Garcia A, Courtney C, Fernandez-De-Las-Penas C, Arroyo-Morales M. Water versus land-based multimodal exercise program effects on body composition in breast cancer survivors: a controlled clinical trial. *Support Care Cancer* 2013; 21: 521–30.
15. Hayes S, Spence R, Galvao D, Newton R. Australian Association for Exercise and Sport Science Position Stand: Optimising cancer outcomes through exercise. *J Sci Med Sport* 2009; 12: 428–34.
16. Courneya KS, McKenzie DC, Mackey JR, et al: Subgroup effects in a randomised trial of different types and doses of exercise during breast cancer chemotherapy. *Br J Cancer* 2014; 111: 1718–1725.
17. Hladiuk M, Huchcroft S, Pemple W, Schurr BE. Arm function after axillary dissection for breast cancer: a pilot study to provide parameter estimates. *J Surg Oncol* 1992; 50: 47–52.
18. Siegel R, DeSantis C, Virgo K, Stein K, Mariotto A et al. Cancer treatment and survivorship statistics, 2012. *CA Cancer J Clin* 2012; 62: 220–24.

19. Skolarus TA, Wolf AM, Erb NL, Brooks DD, Rivers BM et al. American Cancer Society prostate cancer survivorship care guidelines. *CA Cancer J Clin* 2014; 64: 225–49.
20. McTiernan A. Mechanisms linking physical activity with cancer. *Nat Rev Cancer* 2008; 8: 205–11.
21. Friedenreich CM, Neilson HK, Lynch BM. State of the epidemiological evidence on physical activity and cancer prevention. *Eur J Cancer* 2010; 46: 593–604.
22. Young-McCaughan S. Potential for prostate cancer prevention through physical activity. *World J Urol* 2012; 30: 167–79.
23. Leitzmann MF. Physical Activity and Genitourinary Cancer Prevention. In: Courneya KS, Friedenreich CM (Hrsg.) *Physical Activity and Cancer*. Springer Verlag, Berlin, Heidelberg 2011, S. 44.
24. Kushi LH, Doyle C, McCullough M, Rock CL, Demark-Wahnefried W et al. American Cancer Society Guidelines on nutrition and physical activity for cancer prevention: reducing the risk of cancer with healthy food choices and physical activity. *CA Cancer J Clin* 2012; 62: 30–67.
25. Machold S. Evidenz physiotherapeutischer Behandlung bei Prostataktomie-Inkontinenz. *Physioscience* 2006; 2: 106–16.
26. Baumann FT, Zopf EM, Bloch W. Clinical exercise interventions in prostate cancer patients--a systematic review of randomized controlled trials. *Support Care Cancer* 2012; 20: 21–33.
27. Schmitz KH, Courneya KS, Matthews C, Demark-Wahnefried W, Galvão DA et al. American College of Sports Medicine roundtable on exercise guidelines for cancer survivors. *Med Sci Sports Exerc* 2010; 42: 409–26.
28. Baumann FT, Herweg C, Schüle K. Bewegungstherapie und Sport bei unterschiedlichen Krebsentitäten. In: Baumann FT, Schüle K (Hrsg.). *Bewegungstherapie und Sport bei Krebs – Leitfaden für die Praxis*. Deutscher Ärzte-Verlag, Köln 2008, S. 105–200.
29. Wiskemann J, Nies, R., Vandenbergh, D. Leukämien und Lymphome. In: Baumann FT, Jäger E, Bloch W (Hrsg.) *Sport und körperliche Aktivität in der Onkologie*. Berlin, Heidelberg: Springer Verlag 2012; S. 189–205.
30. Wagner LI, Cella D. Fatigue and cancer: causes, prevalence and treatment approaches. *Br J Cancer* 2004; 91: 822–8. doi:10.1038/sj.bjc.6602012.
31. Stubblefield MD, Burstein HJ, Burton AW, Custodio CM, Deng GE et al. NCCN task force report: management of neuropathy in cancer. *J Natl Compr Canc Net* 2009; 7 Suppl 5: S1–S26.; quiz S7–8.
32. Baker KS, Ness KK, Weisdorf D, Francisco L, Sun CL et al. Late effects in survivors of acute leukemia treated with hematopoietic cell transplantation: a report from the Bone Marrow Transplant Survivor Study. *Leukemia* 2010; 24: 2039–47.
33. Friedenreich CM. Physical activity and cancer prevention: from observational to intervention research. *Cancer Epidemiol Biomarkers Prev* 2001; 10: 287–301.
34. Baumann FT, Kraut L, Schule K, Bloch W, Fauser AA. A controlled randomized study examining the effects of exercise therapy on patients undergoing haematopoietic stem cell transplantation. *Bone marrow transplant* 2010; 45: 355–62.
35. Jarden M, Baadsgaard MT, Hovgaard DJ, Boesen E, Adamsen L. A randomized trial on the effect of a multimodal intervention on physical capacity, functional performance and quality of life in adult patients undergoing allogeneic SCT. *Bone marrow transplant* 2009; 43: 725–37.
36. Streckmann F, Kneis S, Leifert JA, Baumann FT, Kleber M et al. Exercise program improves therapy-related side-effects and quality of life in lymphoma patients undergoing therapy. *Ann Oncol* 2014; 25: 493–9.
37. Chang PH, Lai YH, Shun SC, Lin LY, Chen ML et al. Effects of a walking intervention on fatigue-related experiences of hospitalized acute myelogenous leukemia patients undergoing chemotherapy: a randomized controlled trial. *J Pain Symptom Manage* 2008; 35: 524–34.
38. Bergenthal N WA, Streckmann F, Wolkewitz K-D, Monsef I, Engert A et al. Aerobic physical exercise for adult patients with haematological malignancies. *Cochrane Database Syst Rev* 2014. doi:10.1002/14651858.CD009075.
39. Jarden M, Nelausen K, Hovgaard D, Boesen E, Adamsen L. The effect of a multimodal intervention on treatment-related symptoms in patients undergoing hematopoietic stem cell transplantation: a randomized controlled trial. *J Pain Symptom Manage* 2009; 38: 174–90.
40. Streckmann F, Rittweger J, Bloch W, Baumann FT. Bewegungsempfehlungen bei Chemotherapie-induzierter peripherer Polyneuropathie. *B&G* 2014: 179–82.
41. Graf N. Tumorerkrankungen im Kindesalter. In: Sitzmann FC (Hrsg.). *Pädiatrie*. 2. Aufl. Stuttgart: Thieme 2002, S. 484–532.
42. San Juan AF, Wolin K, Lucia A. Physical activity and pediatric cancer survivorship. *Recent Results Cancer Res* 2011; 186: 319–47.
43. Soares-Miranda L, Fuiza-Luces C, Lucia A. Exercise in pediatric cancer patients. In: Ulrich CM, Steindorf K, Berger NA (eds.). *Exercise, energy balance, and cancer*. New York: Springer 2013: 159–79.
44. Oeffinger KC, Mertens AC, Sklar CA, Kawashima T, Hudson MM et al. Chronic health conditions in adult survivors of childhood cancer. *N Engl J Med* 2006; 355: 1572–82.
45. Bhatia S, Meadows AT. Long-term follow-up of childhood cancer survivors: future directions for clinical care and research. *Ped Blood Cancer* 2006; 46: 143–8.

46. Baumann FT, Bloch W, Beulertz J. Clinical exercise interventions in pediatric oncology – A systematic review. *Pediatr Res* 2013; 74: 366–74.
47. Winter C, Müller C, Brandes M, Brinkmann A, Hoffmann C et al. Level of activity in children undergoing cancer treatment. *Pediatr Blood Cancer* 2009; 53: 438–43.
48. Chamorro-Viña C, Ruiz JR, Santana-Sosa E, González VM, Madero L et al. Exercise during hematopoietic stem cell transplant hospitalization in children. *Med Sci Sports Exerc* 2010; 42: 1045–53.
49. Hartman A, te Winkel ML, van Beek RD, de Muinck Keizer-Schrama SM, Kemper HC et al. A randomized trial investigating an exercise program to prevent reduction of bone mineral density and impairment of motor performance during treatment for childhood acute lymphoblastic leukemia. *Pediatr Blood Cancer* 2009; 53: 64–71.
50. Hinds P, Hockenberry M, Rai S, Zhang L, Razzouck BI et al. Clinical field testing of an enhanced activity intervention in hospitalized children with cancer. *J Pain Symptom Manage* 2007; 6: 686–97.
51. Marchese VG, Chiarello LA, Lange BJ. Effects of physical therapy intervention for children with acute lymphoblastic leukemia. *Pediatr Blood Cancer* 2004; 42: 127–33.
52. Rosenhagen A, Bernhörster M, Vogt L, Weiss B, Senn A et al. Implementation of structured physical activity in the pediatric stem cell transplantation. *Klin Pädiatr* 2011; 223: 147–51.
53. San Juan AF, Chamorro-Viña C, Moral S, Fernández del Valle M, Madero C et al. Benefits of intrahospital exercise training after pediatric bone marrow transplantation. *Int J Sports Med* 2008; 29: 439–46.
54. Keats MR, Culos-Reed SN. A community-based physical activity program for adolescents with cancer (project TREK): program feasibility and preliminary findings. *J Pediatr Hematol Oncol* 2008; 30: 272–80.
55. Perondi MB, Gualano B, Artioli GG, de Salles Painelli V, Filho VO et al. Effects of a combined aerobic and strength training program in youth patients with acute lymphoblastic leukemia. *J Sports Sci Med* 2012; 11: 387–92.
56. Ruiz J, Fleck S, Vingren J, Ramírez M, Madero L et al. Preliminary findings of a 4-month intrahospital exercise training intervention on TGFs and TGFβPs in children with leukemia. *J Strength Cond Res* 2010; 24: 1292–7.
57. San Juan AF, Fleck SJ, Chamorro-Viña C, Maté-Munoz JL, Moral S et al. Effects of an intrahospital exercise program intervention for children with leukemia. *Med Sci Sports Exerc* 2007; 39: 13–21.
58. Takken T, van der Torre P, Zwerink M, Hulzebos EH, Bierings M et al. Development, feasibility and efficacy of a community-based exercise training program in pediatric cancer survivors. *Psychooncology* 2009; 18: 440–8.
59. Beulertz J, Bloch W, Prokop A, Baumann FT. Zielgerichtete und individualisierte Bewegungstherapie in der pädiatrischen Onkologie – ein Pilotprojekt. *Monatsschr Kinderheilk* 2013; 4: 330–5.
60. Weiss Kelly AK. Physical activity prescription for childhood cancer survivors. *Curr Sports Med Rep* 2011; 10: 352–9.

Literatur zu Kapitel 26 Organtransplantation

1. http://www.eurotransplant.org/cms/mediaobject.php?file=AR_ET_20153.pdf
2. www.dso.de/organspende-und-transplantation/transplantation.
3. Bussières LM, Pflugfelder PW, Taylor AW, Noble EG, Kostuk WJ. Changes in skeletal muscle morphology and biochemistry after cardiac transplantation. *Am J Cardiol* 1997; 79: 630–634.
4. Mitsui T, Azuma H, Nagasawa M, Iuchi T, Akaike M, et al. Chronic corticosteroid administration causes mitochondrial dysfunction in skeletal muscle. *J Neurol* 2002; 249: 1004–1009.
5. Tegtbu U, Busse MW, Jung K et al. Phase III rehabilitation after heart transplantation. *Z Kardiol* 2003; 92: 908–915.
6. Braith RW, Mills RM, Welsch MA, Keller JW, Pollock ML. Resistance exercise training restores bone mineral density in heart transplant recipients. *J Am Coll Cardiol* 1996; 15: 28: 1471–7.
7. Langer D, Burtin C, Schepers L, Ivanova A, Verleden G, et al. Exercise training after lung transplantation improves participation in daily activity: a randomized controlled trial. *Am J Transplant* 2012; 12: 1584–92.
8. Tegtbu U, Gottlieb J, Bara C, Dierich M, Gützlaff E, et al. Home Based Exercise Training in Lung and Heart Transplant Recipients – Effects on Maximum and Endurance Exercise Capacity. *J Heart Lung Transplant* 2007; 11: 423.
9. Mitchell MJ, Baz MA, Fulton MN, Lisor CF, Braith RW. Resistance training prevents vertebral osteoporosis in lung transplant recipients. *Transplantation* 2003; 15: 76: 557–62.
10. Painter PL, Hector L, Ray K, Lynes L, Dibble S, et al. A randomized trial of exercise training after renal transplantation. *Transplantation* 2002; 74: 42–8.
11. Krasnoff JB, Vintro AQ, Ascher NL, Bass NM, Paul SM, et al. A randomized trial of exercise and dietary counseling after liver transplantation. *Am J Transplant* 2006; 6: 1896–905.

Literatur zu Kapitel 27 Fibromyalgie-Syndrom

1. Reimers CD, Knapp G, Reimers N (2015) Fibromyalgie-Syndrom. In: Reimers CD, Reuter I, Tettenborn B, Mewes N, Knapp G (Hrsg.) Prävention und Therapie durch Sport. Band 2: Neurologie, Psychiatrie/Psychosomatik, Schmerzsyndrome. 2. Aufl., Elsevier Urban & Fischer, München, S. 469–493.
2. Busch AJ, Webber SC, Brachanec M, Bidonde J, Bello-Haas VD et al. Exercise therapy for fibromyalgia. *Curr Pain Headache Rep* 2011; 15: 358–67.
3. Chinn S, Caldwell W, Gritsenko K. Fibromyalgia pathogenesis and treatment options update. *Curr Pain Headache Rep* 2016; 20: 25.
4. Macfarlane GJ, Kronisch C, Dean LE, Atzeni F, Häuser W et al. EULAR revised recommendations for the management of fibromyalgia. *Ann Rheum Dis* 2017; 76: 318–32.
5. Bidonde J, Busch AJ, Bath B, Milosavljevic S. Exercise for adults with fibromyalgia: an umbrella systematic review with synthesis of best evidence. *Curr Rheumatol Rev* 2014; 10: 45–79.
6. Bidonde J, Busch AJ, Webber SC, Schachter CL, Danyliw A et al. Aquatic exercise training for fibromyalgia. *Cochrane Database Syst Rev* 2014; CD011336.
7. Bardal EM, Roeleveld K, Mork PJ. Aerobic and cardiovascular autonomic adaptations to moderate intensity endurance exercise in patients with fibromyalgia. *J Rehabil Med* 2015; 47: 639–46.
8. Larsson A, Palstam A, Löfgren M, Ernberg M, Bjersing J et al. Resistance exercise improves muscle strength, health status and pain intensity in fibromyalgia—a randomized controlled trial. *Arthritis Res Ther* 2015; 17: 161.
9. Busch AJ, Webber SC, Richards RS, Bidonde J, Schachter CL et al. Resistance exercise training for fibromyalgia. *Cochrane Database Syst Rev* 2013; CD010884.
10. Duruturk N, Tuzun EH, Culhaoglu B. Is balance exercise training as effective as aerobic exercise training in fibromyalgia syndrome? *Rheumatol Int* 2015; 35: 845–54.

Literatur zu Kapitel 28 Osteoporose

1. WHO. Osteoporosis – definition and treatment. *Am J Med* 1991; 90: 107–10.
2. DVO. DVO-Leitlinien 2014 zur Prophylaxe, Diagnostik und Therapie der Osteoporose bei Männern ab dem 60. Lebensjahr und postmenopausalen Frauen. Stuttgart: Schattauer; 2014.
3. Moayyeri A. The association between physical activity and osteoporotic fractures: a review of the evidence and implications for future research. *Ann Epidemiol* 2008; 18: 827–35.
4. Kemmler W, Haberle L, von Stengel S. Effects of exercise on fracture reduction in older adults : A systematic review and meta-analysis. *Osteoporos Int* 2013; 24: 1937–50.
5. Kemmler W, Bebenek M, Kohl M, Von Stengel S. Exercise and fractures in postmenopausal women. Final results of the controlled Erlangen Fitness and Osteoporosis Prevention Study (EFOPS). *Osteoporos Int* 2015; 26: 2491–9.
6. Howe TE, Shea B, Dawson LJ, Downie F, Murray A, Ross C, et al. Exercise for preventing and treating osteoporosis in postmenopausal women. *Cochrane Database Syst Rev* 2011: CD000333.
7. Kemmler W, von Stengel S. Exercise and osteoporosis-related fractures: Perspectives and recommendations of the sports and exercise scientist. *Phys Sportsmed* 2011; 39: 142–57.
8. Kemmler W, Weineck J, Hensen J, Lauber D, Kalender WA, Engelke K. Empfehlungen für ein körperliches Training zur Verbesserung der Knochenfestigkeit: Schlussfolgerungen aus Tiermodellen und Untersuchungen an Leistungssportlern. *Dtsch Z Sportmed* 2003; 54: 306–16.
9. Raymond MJ, Bramley-Tzerefos RE, Jeffs KJ, Winter A, Holland AE. Systematic review of high-intensity progressive resistance strength training of the lower limb compared with other intensities of strength training in older adults. *Arch Phys Med Rehabil* 2013; 94: 1458–72.
10. Steib S, Schoene D, Pfeifer K. Dose-response relationship of resistance training in older adults: a meta-analysis. *Med Sci Sports Exerc* 2010; 42: 902–14.
11. Fukumoto Y, Tateuchi H, Ikezoe T, Tsukagoshi R, Akiyama H, So K, et al. Effects of high-velocity resistance training on muscle function, muscle properties, and physical performance in individuals with hip osteoarthritis: a randomized controlled trial. *Clin Rehabil* 2014; 28: 48–58.
12. Slatkovska L, Alibhai SM, Beyene J, Cheung AM. Effect of whole-body vibration on BMD: a systematic review and meta-analysis. *Osteoporos Int* 2010; 21: 1969–80.
13. Burr DB, Robling AG, Turner CH. Effects of biomechanical stress on bones in animals. *Bone* 2002; 30: 781–6.
14. Kemmler W, von Stengel S, Kohl M. Exercise frequency and bone mineral density development in exercising postmenopausal osteopenic women. Is there a critical dose of exercise for affecting bone? Results of the Erlangen Fitness and Osteoporosis Prevention Study. *Bone* 2016; 89: 1–6.

Literatur zu Kapitel 29 Unspezifische Kreuzschmerzen

1. Manchikanti L, Singh V, Falco FJ, Benyamin RM, Hirsch JA. Epidemiology of low back pain in adults. *Neuromodulation* 2014; 17 Suppl 2: 3–10.
2. Mortazavi J, Zebardast J, Mirzashahi B. Low back pain in athletes. *Asian J Sports Med* 2015; 6: e24718.
3. Moradi V, Memari AH, Shayestehfar M, Kordi R. Low back pain in athletes is associated with general and sport specific risk factors: A comprehensive review of longitudinal studies. *Rehabil Res Pract* 2015; 2015: 850184.
4. Reimers N, Reimers CD, Janßen A, Knapp G. Unspezifische Kreuzschmerzen. In: Mooren F, Knapp G, Reimers CD (Hrsg.) *Prävention und Therapie durch Sport. Band 3: Orthopädie, Rheumatologie, Immunologie. 2. Aufl., München: Elsevier Urban & Fischer, 2016, S. 137–78.*
5. Trompeter K, Fett D, Platen P. Prevalence of back pain in sports: A systematic review of the literature. *Sports Med.* 2016 Dec 29. doi: 10.1007/s40279-016-0645-3. [Epub ahead of print]
6. Steffens D, Maher CG, Pereira LS, Stevens ML, Oliveira VC et al. Prevention of low back pain: A systematic review and meta-analysis. *JAMA Intern Med* 2016; 176:199–208.
7. Searle A, Spink M, Ho A, Chuter V. Exercise interventions for the treatment of chronic low back pain: a systematic review and meta-analysis of randomised controlled trials. *Clin Rehabil* 2015; 29: 1155–67.
8. Smith BE, Littlewood C, May S. An update of stabilisation exercises for low back pain: a systematic review with meta-analysis. *BMC Musculoskelet Disord* 2014; 15: 416.
9. Steele J, Bruce-Low S, Smith D. A review of the clinical value of isolated lumbar extension resistance training for chronic low back pain. *PM R* 2015; 7: 169–87.
10. Aladro-Gonzalvo AR, Araya-Vargas GA, Machado-Díaz M, Salazar-Rojas W. Pilates-based exercise for persistent, non-specific low back pain and associated functional disability: a meta-analysis with meta-regression. *J Bodyw Mov Ther* 2013; 17: 125–36.
11. Lin HT, Hung WC, Hung JL, Wu PS, Liaw LJ, Chang JH. Effects of pilates on patients with chronic non-specific low back pain: a systematic review. *J Phys Ther Sci* 2016; 28: 2961–9.
12. Miyamoto GC, Costa LO, Cabral CM. Efficacy of the Pilates method for pain and disability in patients with chronic nonspecific low back pain: a systematic review with meta-analysis. *Braz J Phys Ther* 2013; 17: 517–32.
13. Wells C, Kolt GS, Marshall P, Hill B, Bialocerkowski A. The effectiveness of Pilates exercise in people with chronic low back pain: a systematic review. *PLoS One* 2014; 9: e100402.
14. Holtzman S, Beggs RT. Yoga for chronic low back pain: a meta-analysis of randomized controlled trials. *Pain Res Manag* 2013; 18: 267–72.
15. Wieland LS, Skoetz N, Pilkington K, Vempati R, D'Adamo CR, Berman BM. Yoga treatment for chronic non-specific low back pain. *Cochrane Database Syst Rev.* 2017 Jan 12; 1: CD010671. doi: 10.1002/14651858.CD010671.pub2.

Literatur zu Kapitel 30 Arthrose

1. Orth P, Madry H. Konservative Therapie der Arthrose. *Orthopädie und Unfallchirurgie up2date* 2013; 8: 419–436. doi:10.1007/s00142-008-0481-z.
2. Bennell KL, Dobson F, Hinman RS. Exercise in osteoarthritis: moving from prescription to adherence. *Best Pract Res Clin Rheumatol* 2014; 28: 93–117. doi:10.1016/j.berh.2014.01.009.
3. Johnson VL, Hunter DJ. The epidemiology of osteoarthritis. *Best Pract Res Clin Rheumatol* 2014; 28: 5–15. doi:10.1016/j.berh.2014.01.004.
4. Fernandes L, Hagen KB, Bijlsma JWJ et al. EULAR recommendations for the non-pharmacological core management of hip and knee osteoarthritis. *Ann Rheum Dis* 2013; 72: 1125–35. doi:10.1136/annrheumdis-2012-202745.
5. Bennell KL, Wrigley T V, Hunt MA et al. Update on the role of muscle in the genesis and management of knee osteoarthritis. *Rheum Dis Clin North Am* 2013; 39: 145–76. doi:10.1016/j.rdc.2012.11.003.
6. Hochberg MC, Altman RD, April KT et al. American College of Rheumatology 2012 recommendations for the use of nonpharmacologic and pharmacologic therapies in osteoarthritis of the hand, hip, and knee. *Arthritis Care Res (Hoboken)* 2012; 64: 465–74. <http://www.ncbi.nlm.nih.gov/pubmed/22563589> (accessed 10.1.2015).
7. Pereira D, Peleteiro B, Araújo J et al. The effect of osteoarthritis definition on prevalence and incidence estimates: a systematic review. *Osteoarthritis Cartilage* 2011; 19:1270–85. doi:10.1016/j.joca.2011.08.009
8. Jordan JM, Helmick CG, Renner JB et al. Prevalence of hip symptoms and radiographic and symptomatic hip osteoarthritis in African Americans and Caucasians: the Johnston County Osteoarthritis Project. *J Rheumatol* 2009; 36: 809–15. doi:10.3899/jrheum.080677.
9. Zhang W, Moskowitz RW, Nuki G et al. OARSI recommendations for the management of hip and knee osteoarthritis, Part II: OARSI evidence-based, expert consensus guidelines. *Osteoarthritis Cartilage* 2008; 16: 137–62. doi:10.1016/j.joca.2007.12.013.

10. RKI. Gesundheitsberichterstattung des Bundes. 2012. doi:978-3-89606-170-6.
11. van Baar ME, Dekker J, Oostendorp R et al. Effectiveness of exercise in patients with osteoarthritis of hip or knee: nine months' follow up. *Ann Rheum Dis* 2001; 60: 1123–30.
12. Felson DT, Lawrence RC, Hochberg MC et al. Osteoarthritis: new insights. Part 2: treatment approaches. *Ann Intern Med* 2000; 133: 726–37. <http://www.ncbi.nlm.nih.gov/pubmed/11074906> (accessed 28.11.2015).
13. Musumeci G, Castrogiovanni P, Trovato FM et al. Physical activity ameliorates cartilage degeneration in a rat model of aging: a study on lubricin expression. *Scand J Med Sci Sports* 2015; 25: e222–30. doi:10.1111/sms.12290.
14. Mayer F, Schmitt H, Dickhuth H-H. Die Bedeutung von Sport in der Entstehung, Prävention und Rehabilitation der Arthrose. *Aktuelle Rheumatol* 2003; 28: 210–7. doi:10.1055/s-2003-41642.
15. Vincent KR, Vincent HK. Resistance Exercise for Knee Osteoarthritis. *PM R* 2013; 4: 45–52. doi:10.1016/j.pmrj.2012.01.019. Resistance.
16. Mayer F, Scharhag-Rosenberger F, Carlsohn A, et al. The intensity and effects of strength training in the elderly. *Dtsch Arztebl Int* 2011; 108: 359–64. doi:10.3238/arztebl.2011.0359.
17. Slemenda C, Heilman DK, Brandt KD, et al. Reduced quadriceps strength relative to body weight: a risk factor for knee osteoarthritis in women? *Arthritis Rheum* 1998; 41: 1951–9. doi:10.1002/1529-0131.
18. Buckwalter JA, Lane NE. Athletics and osteoarthritis. *Am J Sports Med*; 25: 873–81. <http://www.ncbi.nlm.nih.gov/pubmed/9397280> (accessed 7.11.2015).
19. Dvorak J. Osteoarthritis in football: FIFA/F-MARC approach. *Br J Sports Med* 2011; 45: 673–6. doi:10.1136/bjsm.2011.083584.
20. Tveit M, Rosengren BE, Nilsson J et al. Former Male Elite Athletes Have a Higher Prevalence of Osteoarthritis and Arthroplasty in the Hip and Knee Than Expected. *Am J Sports Med* 2012; 40: 527–33. doi:10.1177/0363546511429278.
21. Zhang Y. Prevalence of symptomatic osteoarthritis and its impact on functional status among the elderly: The framingham study. *Am J Epidemiol* 2002; 156: 1021–7. doi:10.1093/aje/kwf141.
22. Manninen P, Riihimäki H, Heliövaara M, et al. Physical exercise and risk of severe knee osteoarthritis requiring arthroplasty. *Rheumatology (Oxford)* 2001; 40: 432–7. <http://www.ncbi.nlm.nih.gov/pubmed/11312383> (accessed 1.12.2015).
23. Williams PT. Effects of running and walking on osteoarthritis and hip replacement risk. *Med Sci Sports Exerc* 2013; 45: 1292–7. doi:10.1249/MSS.0b013e3182885f26.
24. Rajabi R, Johnson GM, Alizadeh MH et al. Radiographic knee osteoarthritis in ex-elite table tennis players. *BMC Musculoskelet Disord* 2012; 13: 12. doi:10.1186/1471-2474-13-12.
25. Michaëlsson K, Byberg L, Ahlborn A et al. Risk of severe knee and hip osteoarthritis in relation to level of physical exercise: A prospective cohort study of long-distance Skiers in Sweden. *PLoS One* 2011; 6: 1–8. doi:10.1371/journal.pone.0018339.
26. Golightly YM, Marshall SW, Callahan LF et al. Early-onset arthritis in retired National Football League players. *J Phys Act Health* 2009; 6: 638–43. <http://www.ncbi.nlm.nih.gov/pubmed/19953841> (accessed 22.10.2016).
27. Schmitt H, Rohs C, Schneider S et al. Is competitive running associated with osteoarthritis of the hip or the knee? *Orthopäde* 2006; 35: 1087–92. doi:10.1007/s00132-006-0998-7.
28. Maquirriain J, Ghisi JP, Amato S. Is tennis a predisposing factor for degenerative shoulder disease? A controlled study in former elite players. *Br J Sports Med* 2006; 40: 447–50. doi:10.1136/bjsm.2005.023382.
29. L'Hermette M, Polle G, Tourny-Chollet C et al. Hip passive range of motion and frequency of radiographic hip osteoarthritis in former elite handball players. *Br J Sports Med* 2006; 40: 45-49-49. doi:10.1136/bjsm.2005.019026.
30. Pedersen BK, Saltin B. Exercise as medicine – Evidence for prescribing exercise as therapy in 26 different chronic diseases. *Scand J Med Sci Sport* 2015; 25: 1–72. doi:10.1111/sms.12581.
31. Zhang W, Nuki G, Moskowitz RW et al. OARSI recommendations for the management of hip and knee osteoarthritis Part III: changes in evidence following systematic cumulative update of research published through January 2009. *YJOCA* 2010; 18: 476–99. doi:10.1016/j.joca.2010.01.013.
32. Pietschmann MF, Wohlleb L, Weber P et al. Sports activities after medial unicompartmental knee arthroplasty Oxford III – What can we expect? *Int Orthop* 2013; 37: 31–7. doi:10.1007/s00264-012-1710-7.
33. Regnaud J, Lefevre-Colau M, Trinquart L et al. High-intensity versus low-intensity physical activity or exercise in people with hip or knee osteoarthritis. *Cochrane database Syst Rev Published Online First*: 2015. doi:10.1002/14651858.CD010203.pub2. www.cochranelibrary.com.
34. Kristensen J, Franklyn-Miller A. Resistance training in musculoskeletal rehabilitation : a systematic review. *Br J Sports Med* 2012; 46: 719–26. doi:10.1136/bjsports79376.
35. Cassel M, Hotzkow K, Mayer F. Arthrosepatienten: Empfehlungen zur sportlichen Belastung. *Ärztliches J Orthopädie/ Rheumatologie* 2013; 4: 6–8.
36. Kise NJ, Risberg MA, Stensrud S et al. Exercise therapy versus arthroscopic partial meniscectomy for degenerative meniscal tear in middle aged patients : randomised controlled trial with two year follow-up. doi:10.1136/bmj.i3740.

37. Gür H, Çakın N, Akova B, et al. Concentric versus combined concentric-eccentric isokinetic training: Effects on functional capacity and symptoms in patients with osteoarthritis of the knee. *Arch Phys Med Rehabil* 2002; 83: 308–16. doi:10.1053/apmr.2002.30620.
38. Bruhn S, Kullmann N, Gollhofer A. Combinatory Effects of High-Intensity-Strength Training and Sensorimotor Training on Muscle Strength. *Int J Sports Med* 2006; 27: 401–6. doi:10.1055/s-2005-865750.
39. Gruber M, Gollhofer A. Impact of sensorimotor training on the rate of force development and neural activation. *Eur J Appl Physiol* 2004; 92: 98–105. doi:10.1007/s00421-004-1080-y.
40. Mayer F, Weber J, Cassel M. Konservative Orthopädie in der Sportmedizin. *Orthopädie und Unfallchirurgie up2date* 2011; 6: 479–98. doi:10.1055/s-0030-1257083.

Literatur zu Kapitel 31 Fatigue

1. Reimers CD (2015) Fatigue. In: Reimers CD, Reuter I, Tettenborn B, Mewes N, Knapp G (Hrsg.) *Prävention und Therapie durch Sport. Band 2: Neurologie, Psychiatrie/Psychosomatik, Schmerzsyndrome*. 2. Aufl., Elsevier Urban & Fischer, München, S. 543–553.
2. Cramer H, Lauche R, Azizi H, Dobos G, Langhorst J. Yoga for multiple sclerosis: a systematic review and meta-analysis. *PLoS One* 2014 Nov 12; 9: e112414. doi: 10.1371/journal.pone.0112414. eCollection 2014.
3. Cramer H, Lauche R, Klose P, Dobos G, Langhorst J. A systematic review and meta-analysis of exercise interventions for colorectal cancer patients. *Eur J Cancer Care* 2014 Jan; 23: 3–14. doi: 10.1111/ecc.12093. Epub 2013 Jul 8.
4. Cramp F, Hewlett S, Almeida C, Kirwan JR, Choy EH, Chalder T, Pollock J, Christensen R. Non-pharmacological interventions for fatigue in rheumatoid arthritis. *Cochrane Database Syst Rev* 2013 Aug 23; 8: CD008322. doi: 10.1002/14651858.CD008322.pub2.
5. Elbers RG, Verhoef J, van Wegen EE, Berendse HW, Kwakkel G. Interventions for fatigue in Parkinson's disease. *Cochrane Database Syst Rev* 2015 Oct 8; 10: CD010925. doi: 10.1002/14651858.CD010925.pub2.
6. Furmaniak AC, Menig M, Markes MH. Exercise for women receiving adjuvant therapy for breast cancer. *Cochrane Database Syst Rev* 2016 Sep 21; 9: CD005001. [Epub ahead of print].
7. Heine M, van de Port I, Rietberg MB, van Wegen EE, Kwakkel G. Exercise therapy for fatigue in multiple sclerosis. *Cochrane Database Syst Rev* 2015 Sep 11: CD009956. doi: 10.1002/14651858.CD009956.pub2.
8. Larun L, Brurberg KG, Odgaard-Jensen J, Price JR. Exercise therapy for chronic fatigue syndrome. *Cochrane Database Syst Rev*. 2016 Jun 24:CD003200. doi: 10.1002/14651858.CD003200.pub5.
9. Marcellis R, Van der Veeke M, Mesters I, Drent M, De Bie R, De Vries G, Lenssen A. Does physical training reduce fatigue in sarcoidosis? *Sarcoidosis Vasc Diffuse Lung Dis* 2015 Jun 22; 32: 53–62.
10. Meneses-Echávez JF, González-Jiménez E, Ramírez-Vélez R. Effects of supervised exercise on cancer-related fatigue in breast cancer survivors: a systematic review and meta-analysis. *BMC Cancer* 2015 Feb 21; 15: 77. doi: 10.1186/s12885-015-1069-4.
11. Meneses-Echávez JF, González-Jiménez E, Ramírez-Vélez R. Supervised exercise reduces cancer-related fatigue: a systematic review. *J Physiother* 2015a Jan; 61: 3–9. doi: 10.1016/j.jphys.2014.08.019. Epub 2014 Dec 13.
12. Puetz TW, Herring MP. Differential effects of exercise on cancer-related fatigue during and following treatment: a meta-analysis. *Am J Prev Med* 2012 Aug; 43: e1–24. doi: 10.1016/j.amepre.2012.04.027.
13. Smith BA, Neidig JL, Nickel JT, Mitchell GL, Para MF, Fass RJ. Aerobic exercise: effects on parameters related to fatigue, dyspnea, weight and body composition in HIV-infected adults. *AIDS* 2001 Apr 13; 15: 693–701.
14. van Vulpen JK, Peeters PH, Velthuis MJ, van der Wall E, May AM. Effects of physical exercise during adjuvant breast cancer treatment on physical and psychosocial dimensions of cancer-related fatigue: A meta-analysis. *Maturitas* 2016 Mar; 85: 104–11. doi: 10.1016/j.maturitas.2015.12.007. Epub 2015 Dec 28.

Literatur zu Kapitel 32 Schlaganfall

1. Reimers CD, Wingendorf I, Holzgraefe M, Reimers AK, Knapp G. Schlaganfälle. In: Reimers CD, Reuter I, Tettenborn B, Mewes N, Knapp G (Hrsg.) *Prävention und Therapie durch Sport. Band 2: Neurologie, Psychiatrie/Psychosomatik, Schmerzsyndrome*. 2. Aufl., München: Elsevier Urban & Fischer, 2015, S. 19–76.
2. Chen X, Zhou L, Zhang Y, Yi D, Liu L et al. Risk factors of stroke in Western and Asian countries: a systematic review and meta-analysis of prospective cohort studies. *BMC Public Health* 2014; 14: 776.
3. Pang MY, Charlesworth SA, Lau RW, Chung RC. Using aerobic exercise to improve health outcomes and quality of life in stroke: evidence-based exercise prescription recommendations. *Cerebrovasc Dis* 2013; 35: 7–22.
4. Kristensen OH, Stenager E, Dalgas U. Muscle strength and poststroke hemiplegia: A systematic review of muscle strength assessment and muscle strength impairment. *Arch Phys Med Rehabil* 2017; 98: 368–80.

5. Kyu HH, Bachman VF, Alexander LT, Mumford JE, Afshin A et al. Physical activity and risk of breast cancer, colon cancer, diabetes, ischemic heart disease, and ischemic stroke events: systematic review and dose-response meta-analysis for the Global Burden of Disease Study 2013. *BMJ* 2016; 354: i3857.
6. Tumaszk MT, Trócoli T, de Oliveira MF, Campos RR, Botelho RV. Do physically active patients have better functional outcome after stroke? A systematic review. *J Stroke Cerebrovasc Dis* 2016; 25: 527–32.
7. Wendel-Vos GCW, Schuit AJ, Feskens EJM, Boshuizen HC, Verschuren WMM et al. Physical activity and stroke. A metaanalysis of observational data. *Int J Epidemiol* 2004; 33: 787–98.
8. Chiuve SE, Rexrode KM, Spiegelman D, Logroscino G, Manson JE, Rimm EB. Primary prevention of stroke by healthy lifestyle. *Circulation* 2008; 118: 947–54.
9. Zhang Y, Tuomilehto J, Jousilahti P, Wang Y, Antikainen R, Hu G. Lifestyle factors on the risks of ischemic and hemorrhagic stroke. *Arch Intern Med* 2011; 171: 1811–8.
10. Billinger SA, Arena R, Bernhardt J, Eng JJ, Franklin BA et al. Physical activity and exercise recommendations for stroke survivors: a statement for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke* 2014; 45: 2532–53.
11. Kono Y, Yamada S, Kamisaka K, Araki A, Fujioka Y et al. Recurrence risk after noncardioembolic mild ischemic stroke in a Japanese population. *Cerebrovasc Dis* 2011; 31: 365–72.
12. Turan TN, Nizam A, Lynn MJ, Egan BM, Le NA et al. Relationship between risk factor control and vascular events in the SAMMPRIS trial. *Neurology* 2017; 88: 379–85.
13. Naci H, Ioannidis JP. Comparative effectiveness of exercise and drug interventions on mortality outcomes: metaepidemiological study. *BMJ* 2013; 347: ff5577.
14. Saunders DH, Sanderson M, Hayes S, Kilrane M, Greig CA et al. Physical fitness training for stroke patients. *Cochrane Database Syst Rev* 2016; 3: CD003316.
15. Pang MY, Charlesworth SA, Lau RW, Chung RC. Using aerobic exercise to improve health outcomes and quality of life in stroke: evidence-based exercise prescription recommendations. *Cerebrovasc Dis* 2013; 35: 7–22.
16. Kendall BJ, Gothe NP. Effect of Aerobic Exercise Interventions on Mobility among Stroke Patients: A Systematic Review. *Am J Phys Med Rehabil* 2016; 95: 214–24.
17. Wist S, Clivaz J, Sattelmayer M. Muscle strengthening for hemiparesis after stroke: A meta-analysis. *Ann Phys Rehabil Med* 2016; 59: 114–24.
18. Chen BL, Guo JB, Liu MS, Li X, Zou J et al. Effect of traditional Chinese exercise on gait and balance for stroke: A systematic review and meta-analysis. *PLoS One* 2015; 10: e0135932.
19. Chen L, Chen J, Peng Q, Chen J, Zou Y, Liu G. Effect of sling exercise training on balance in patients with stroke: A meta-analysis. *PLoS One* 2016; 11: e0163351.
20. Stoller O, de Bruin ED, Knols RH, Hunt KJ. Effects of cardiovascular exercise early after stroke: systematic review and metaanalysis. *BMC Neurology* 2012; 12: 45.
21. Brazzelli M, Saunders DH, Greig CA, Mead GE. Physical fitness training for patients with stroke: updated review. *Stroke* 2012; 43: e39–e40.
22. Mijajlović MD, Pavlović A, Brainin M, Heiss WD, Quinn TJ et al. Post-stroke dementia – a comprehensive review. *BMC Med* 2017; 15: 11.
23. Barak S, Hutzler Y, Dubnov-Razi G. Physical exercise after stroke. Effects, recommendations and barriers. *Harefuah* 2016; 155: 378–384.
24. Thayabaranathan T, Andrew NE, Immink MA, Hillier S, Stevens P et al. Determining the potential benefits of yoga in chronic stroke care: a systematic review and meta-analysis. *Top Stroke Rehabil* 2017: 1–10.

Literatur zu Kapitel 33 Parkinson-Syndrom

1. Oertel W, Deuschl G, Poewe W. Parkinson-Syndrome und andere Bewegungsstörungen. Stuttgart: Thieme Verlag 2012.
2. Hu G, Jousilahti P, Nissinen A, Antikainen R, Kivipelto M, Tuomilehto J. Body mass index and the risk of Parkinson's disease. *Neurology* 2006; 67: 1955–9.
3. Sascio AJ, Paffenbarger RS Jr, Gendle I, Wing AL. The role of physical exercise in the occurrence of Parkinson's disease. *Arch Neurol* 1992; 49: 360–5.
4. Xu Q, Park Y, Huang X, Hollenbeck A, Blair A et al. Physical activities and future risk of Parkinson disease. *Neurology* 2010; 75: 341–8.
5. Thacker EL, Chen H, Patel AV, McCullough ML, Calle EE et al. Recreational physical activity and risk of Parkinson's disease. *Mov Disord* 2008; 23: 69–74.
6. Chen H, Zhang SM, Schwarzschild MA, Herman MA, Ascherio A. Physical activity and the risk of Parkinson's disease. *Neurology* 2005; 64: 664–9.

7. Logroscino G, Sesso HD, Paffenberger HS, Lee IM. Physical activity and risk of Parkinson's disease: a prospective cohort study. *J Neurol Neurosurg Psychiatry* 2006; 77: 1318–22.
8. Hamer M, Chida Y. Physical activity and risk of neurodegenerative disease: a systematic review of prospective evidence. *Psychol Med* 2009; 39: 3–11.
9. Kuroda K, Tatara K, Takatorige T, Shinsho F. Effect of physical exercise on mortality in patients with Parkinson's disease. *Acta Neurol Scand* 1992; 86: 55–9.
10. Ahlskog JE. Does vigorous exercise have a neuroprotective effect in Parkinson's disease? *Neurology* 2011; 77: 288–94.
11. Fisher BE, Wu AD, Salem GJ, Song J, Lin CH et al. The effect of exercise training in improving motor performance and corticomotor excitability in people with early Parkinson's disease. *Arch Phys Med Rehabil* 2008; 89: 1221–9.
12. Tanaka K, De Quadros AC, Santos RF, Stella F, Gobbi LTB, Gobbi S. Benefits of physical exercise on executive functions on older people with Parkinson's disease. *Brain Cogn* 2009; 69: 435–41.
13. Cruise KE, Bucks RS, Loftus AM, Newton RU, Pegoraro R, Thomas MG. Exercise and Parkinson's: benefits for cognition and quality of life. *Acta Neurol Scand* 2011; 123: 13–9.
14. Uc EY, Doerschug KC, Magnotta V, Dawson JD, Thomsen TR et al. Phase I/II randomized trial of aerobic exercise in Parkinson's disease in a community setting. *Neurology* 2014; 83: 413–25.
15. Sunwoo MK, Lee JE, Hong JY, Ye BS, Lee HS et al. Premorbid exercise engagement and motor reserve in Parkinson's disease. *Parkinsonism Relat Disord* 2017; 34: 49–53.
16. Koller WC, Kase S. Muscle strength testing in Parkinson's disease. *Eur Neurol* 1986; 25: 130–3.
17. Bridgewater KJ, Sharpe MH. Trunk muscle training and early Parkinson's Disease. *Physiother Theor Pract* 1997; 13: 139–53.
18. Pedersen SW, Oberg B. Dynamic strength in Parkinson's disease. *Eur Neurol* 1993; 33: 97–102.
19. Hass CJ, Bishop M, Moscovich M, Stegemöller EL, Skinner J et al. Defining the clinically meaningful difference in gait speed in persons with Parkinson's disease. *J Neurol Phys Ther* 2014; 38: 233–8.
20. Ebersbach G, Ebersbach A, Edler D, Kauffhold O, Kusch M et al. Comparing exercise in Parkinson's disease-The Berlin LSVT® BIG Study. *Mov Dis* 2010; 25: 1902–08.
21. Reuter I, Mehnert S, Leone P, Kaps M, Oechsner M, Engelhardt M. Effects of Gymnastics, Walking and Nordic Walking in Parkinson's disease. *J Aging Res* 2011; 2011: 232473.
22. Jöbges M, Spittler-Schneiders H, Renner CI, Hummelsheim H. Clinical relevance of rehabilitation programs for Parkinson's disease. I: Non-symptom-specific therapeutic approaches. *Parkinsonism Relat Disord* 2007; 13: 195–202.
23. Jöbges EM, Spittler-Schneiders H, Renner CI, Hummelsheim H. Clinical relevance of rehabilitation programs for patients with idiopathic Parkinson syndrome. II: Symptom-specific therapeutic approaches. *Parkinsonism Relat Disord* 2007; 13: 203–13.
24. Schenkman M, Hall DA, Barón AE, Schwartz RS, Mettler P, Kohrt WM. Exercise for people in early- or mid-stage Parkinson disease: a 16-month randomized controlled trial. *Phys Ther* 2012; 92: 1395–410.
25. Li F, Harmer P, Liu Y, Eckstrom E, Fitzgerald K et al. A randomized controlled trial of patient-reported outcomes with tai chi exercise in Parkinson's disease. *Mov Disord* 2014; 29: 539–45.
26. Leavy B, Roaldsen KS, Nylund K, Hagströmer M, Franzén E. "Pushing the Limits": Rethinking motor and cognitive resources after a highly challenging balance training program for Parkinson disease. *Phys Ther* 2016 Aug 11. [Epub ahead of print].
27. Conradsson D, Löfgren N, Nero H, Hagströmer M, Ståhle A et al. The effects of highly challenging balance training in elderly with Parkinson's disease: A randomized controlled trial. *Neurorehabil Neural Repair* 2015; 29: 827–36.
28. Allen NE, Sherrington C, Paul SS, Canning CG. Balance and falls in Parkinson's disease: a metaanalysis of the effect of exercise and motor training. *Mov Disord* 2011; 26: 1605–15.
29. Toole T, Maitland CG, Warren E, Hubmann MF, Panton L. The effects of loading and unloading treadmill walking on balance, gait, fall risk, and daily function in Parkinsonism. *NeuroRehabilitation* 2005; 20: 307–22.
30. Toole T, Hirsch MA, Forkink A, Lehman DA, Maitland CG. The effect of a balance and strength training program on Parkinsonism: a preliminary study. *J Neurol Rehabil* 2012; 39: 236–47.
31. Hirsch M, Toole T, Maitland CG, Rider RA. The effects of balance training and high-intensity resistance training on persons with idiopathic Parkinson's disease. *Arch Phys Med Rehabil* 2003; 84: 1109–17.
32. Dibble LE, Hale TF, Marcus RL, Gerber JP, La Stayo PC. High intensity eccentric resistance training decreases bradykinesia and improves quality of life in persons with Parkinson's disease: A preliminary study. *Parkinsonism and Relat Disord* 2009; 15: 752–7.
33. Christiansen CL, Schenkman ML, McFann K, Wolfe P, Kohrt WM. Walking economy in people with Parkinson's disease. *Mov Disord* 2009; 24: 1481–7.
34. Eijkeren van FJ, Reijmers RS, Kleinveld MJ, Minten A, Brugen JP, Bloem BR. Nordic walking improves mobility in Parkinson's disease. *Mov Dis* 2008; 23: 2239–49.

35. Herman T, Giladi N, Hausdorff JM. Treadmill training for the treatment of gait disturbances in people with Parkinson's disease; a mini-review. *J Neural Transm* 2009; 116: 307–18.
36. Mehrholz J, Kugler J, Storch A, Pohl M, Hirsch K, Elsner B. Treadmill training for patients with Parkinson's disease. *Cochrane Database Syst Rev* 2015; 13: CD007830.
37. Hass CJ, Malczak P, Nocera J, Stegemöller EL, Wagle Shukla A et al. Quantitative normative gait data in a large cohort of ambulatory persons with Parkinson's disease. *PLoS One* 2012; 7: e42337.
38. Ashburn A, Fazakarley L, Ballinger C, Pickering R, McLellan LD, Fitton C. A randomised controlled trial of a home-based exercise programme to reduce risk of falling among people with Parkinson's disease. *J Neurol Neurosurg Psychiatry* 2007; 78: 678–84.
39. Schmitz-Hübsch T, Derek P, Kielwein K, Fimmers R, Klockgether T, Wüllner U. Qigong exercise for the symptoms of Parkinson's disease: A randomized, controlled pilot study. *Mov Disord* 2006; 21: 543–8.
40. Fritz B, Rombach S, Godau J, Berg D, Horstmann T, Grau S. The influence of Nordic Walking training on sit-to-stand transfer in Parkinson patients. *Gait Posture* 2011; 34: 234–8.
41. Eijkeren van FJ, Reijmers RS, Kleinveld MJ, Minten A, Brugen JP, Bloem BR. Nordic walking improves mobility in Parkinson's disease. *Mov Dis* 2008; 23: 2239–49.
42. Bergen BL, Toole T, Elliott RG, Wallace B, Robinson K, Maitland CG. Aerobic exercise intervention improves aerobic capacity and movement initiation in Parkinson's disease patients. *NeuroRehabilitation* 2002; 17: 161–8.
43. Shen X, Wong-Yu IS, Mak MK. Effects of exercise on falls, balance, and gait ability in Parkinson's disease: A meta-analysis. *Neurorehabil Neural Repair* 2016; 30: 512–27.
44. Cakit BD, Saracoglu M, Genc H, Erdem HR, Inan L. The effects of incremental speed-dependent treadmill training on postural instability and fear of falling in Parkinson's disease. *Clin Rehabil* 2007; 21: 698–705.
45. Ellis T, de Goede CJ, Feldman RG, Wolters EC, Kwakkel G, Wagenaar RC. Efficacy of a physical therapy program in patients with Parkinson's disease: a randomised controlled trial. *Arch Phys Med Rehabil* 2005; 86: 626–32.
46. Miyai I, Fujimoto Y, Yamamoto H, Ueda Y, Saito T et al. Long-term effect of body weight-supported treadmill training in Parkinson's disease: a randomized controlled trial. *Arch Phys Med Rehabil* 2002; 83: 370–3.
47. Hass CJ, Collins MA, Juncos JL. Resistance training with creatinine monohydrate improves upper-body strength in patients with Parkinson's disease: A randomized trial. *Neurorehabil Neural Repair* 2007; 21: 107–15.
48. Schilling BK, Pfeiffer RF, LeDoux MS, Karlage RE, Bloomer RJ, Falvo MJ. Effects of moderate-volume, high-load lower-body resistance training on strength and function in persons with Parkinson's disease: A pilot study. *Parkinsons Dis* 2010; 824734.
49. Chung CL, Thilarajah S, Tan D. Effectiveness of resistance training on muscle strength and physical function in people with Parkinson's disease: a systematic review and meta-analysis. *Clin Rehabil* 2016; 30: 11–23.
50. Protas EJ, Mitchell K, Williams A, Qureshy H, Caroline K, Lai EC. Gait and step training to reduce falls in Parkinson's disease. *NeuroRehabilitation* 2005; 20: 183–90.
51. Schenkman M, Cutson TM, Kuchibhatla M, Chandler JM, Pieper KF et al. Exercise to improve spinal flexibility and function for people with Parkinson's disease: a randomised controlled trial. *J Am Geriatr Soc* 1998; 4: 1207–16.
52. Pohl M, Rockstroh K, Ruckriem S, Mrass G, Mehrholz I. Immediate effects of speed-dependent treadmill training on gait parameters in early Parkinson's disease. *Arch Phys Med Rehabil* 2003; 84: 1760–6.
53. de Dreu MJ, van der Wilk AS, Poppe E, Kwakkel G, van Wegen EE. Rehabilitation, exercise therapy and music in patients with Parkinson's disease: a meta-analysis of the effects of music-based movement therapy on walking ability, balance and quality of life. *Parkinsonism Relat Disord* 2012; 18, Suppl 1: S114–9.
54. Sharp K, Hewitt J. Dance as an intervention for people with Parkinson's disease: a systematic review and meta-analysis. *Neurosci Biobehav Rev* 2014; 47: 445–56.
55. Hackney ME, Earhart GM. Effects of dance on movement control in Parkinson's disease: a comparison of Argentine tango and American ballroom. *J Rehabil Med* 2009; 41: 475–81.
56. Hackney ME, Earhart GM. Short duration, intensive tango dancing for Parkinson disease: an uncontrolled pilot study. *Complement Ther Med* 2009; 17: 203–7. doi: 10.1016/j.ctim.2008.10.005.
57. Rafferty MR, Schmidt PN, Luo ST, Li K, Marras C et al. Regular exercise, quality of life, and mobility in Parkinson's disease: A longitudinal analysis of National Parkinson Foundation Quality Improvement Initiative Data. *J Parkinsons Dis* 2017; 7: 193–202.
58. Cohen AD, Tillerson JL, Smith AD, Schallert T, Zigmond MJ. Neuroprotective effects of prior limb use in 6-hydroxydopamine treated rats: Possible role of GDNF. *J Neurochem* 2003; 85: 299–305.
59. Vučković MG, Fisher B, Nacca A, Leahy RM, Walsh JP et al. Exercise elevates dopamine D2 receptor in a mouse model of Parkinson's disease: in vivo imaging with [¹⁸F] Raclopride. *Mov Dis* 2010; 25: 2777–84.

Literatur zu Kapitel 34 Restless-Legs-Syndrom

1. Aliasgharpour M, Abbasi Z, Pedram Razi S, Kazemnezhad A. The Effect of Stretching Exercises on Severity of Restless Legs Syndrome in Patients on Hemodialysis. *Asian J Sports Med* 2016; 7: e31001. doi: 10.5812/asjrm.31001. eCollection 2016.
2. Batool-Anwar S, Li Y, De Vito K, Malhotra A, Winkelmann J, Gao X. Lifestyle Factors and Risk of Restless Legs Syndrome: Prospective Cohort Study. *J Clin Sleep Med* 2016; 12: 187–94. doi: 10.5664/jcsm.5482.
3. Schlesinger I, Erikh I, Avizohar O, Sprecher E, Yarnitsky D. Cardiovascular risk factors in restless legs syndrome. *Mov Disord* 2009; 4: 587–92. doi: 10.1002/mds.22486.
4. Merlino G, Gigli GL. Sleep-related movement disorders. *Neurol Sci* 2012; 33: 491–513.
5. Phillips B, Young T, Finn L, Asher K, Hening WA, Purvis C. Epidemiology of restless legs symptoms in adults. *Arch Intern Med* 2000; 160: 2137–2141.
6. Reimers CD, Knapp G. Restless-legs-Syndrom und periodische Beinbewegungen. In: Reimers CD, Reuter I, Tettenborn B, Broocks A, Thürauf N, Knapp G (Hrsg.) *Prävention und Therapie durch Sport*. Band 2: Neurologie, Psychiatrie/ Psychosomatik, Schmerzsyndrome. 2. Aufl., Urban & Fischer, München 2015, S. 103–111.
7. Simon J-P, Debout C, Bertran F, Kerouanton A, Schaeffer S, Chapon F. Restless legs syndrome associated with exercise intolerance: Data from a retrospective observational clinical neuromuscular center study. *Rev Neurol* 2013; 169: 663–669.

Literatur zu Kapitel 35 Epilepsie

1. Eom S, Lee MK, Park J-H, Jeon JY, Kang H-C, Lee JS et al. The impact of an exercise therapy on psychosocial health of children with benign epilepsy: a pilot study. *Epilepsy Behav* 2014; 37: 151–156.
2. Kamel JT, Badawy RAB, Cook MJ. Exercise-induced seizures and lateral asymmetry in patients with temporal lobe epilepsy. *Epilepsy Behav Case Rep* 2014; 2: 26–30.
3. Kim J, Jung DS, Hwang KJ, Seo J-H, Na G-Y, Hong SB et al. Can an exercise bicycle be safely used in the epilepsy monitoring unit?: An exercise method to provoke epileptic seizures and the related safety issues. *Epilepsy Behav* 2015; 46: 79–83.
4. Capovilla G, Kaufman KR, Perucca E, Moshé SL, Arida RM. Epilepsy, seizures, physical exercise, and sports: A report from the ILAE Task Force on Sports and Epilepsy. *Epilepsia* 2016; 57: 6–12.

Literatur zu Kapitel 36 Multiple Sklerose

1. Latimer-Cheung AE, Pilutti LA, Hicks AL, Martin Ginis KA, Fenuta AM, MacKibbin KA et al. Effects of exercise training on fitness, mobility, fatigue, and health-related quality of life among adults with multiple sclerosis: a systematic review to inform guideline development. *Arch Phys Med Rehabil* 2013; 94: 1800–28.e3.
2. Ortiz-Rubio A, Cabrera-Martos I, Rodriguez-Torres J, Fajardo-Contreras W, Diaz-Pelegrina A, Valenza MC. Effects of a Home-Based Upper Limb Training Program in Patients With Multiple Sclerosis: A Randomized Controlled Trial. *Arch Phys Med Rehabil* 2016; 97: 2027–33.
3. Kucuk F, Kara B, Poyraz EC, Idiman E. Improvements in cognition, quality of life, and physical performance with clinical Pilates in multiple sclerosis: a randomized controlled trial. *Journal of physical therapy science* 2016; 28: 761–8.
4. Coote S, Hughes L, Rainsford G, Minogue C, Donnelly A. Pilot randomized trial of progressive resistance exercise augmented by neuromuscular electrical stimulation for people with multiple sclerosis who use walking aids. *Arch Phys Med Rehabil* 2015; 96: 197–204.
5. Kjølhede T, Vissing K, de Place L, Pedersen BG, Ringgaard S, Stenager E et al. Neuromuscular adaptations to long-term progressive resistance training translates to improved functional capacity for people with multiple sclerosis and is maintained at follow-up. *Mult Scler* 2015; 21: 599–611.
6. Siemonsen S SJ, Kjølhede T, Wenzel D, Ringgaard S, Pedersen BG, Stenager E, Petersen T, Heesen C, Vissing K, Dalgas U. Resistance training increases cortical thickness in RRMS – results of a pilot RCT. 2015.
7. Hodecker SC SS, Kjølhede T, Ringgaard S, Pedersen BG, Stenager E, Petersen T, Heesen C, Vissing K, Dalgas U, Stellmann JP. Physical exercise-related changes in structural connectome architecture in patients with relapsing-remitting multiple sclerosis. *Multiple Sclerosis Journal* 2016; 22: 7–87.
8. Medina-Perez C, de Souza-Teixeira F, Fernandez-Gonzalo R, de Paz-Fernandez JA. Effects of a resistance training program and subsequent detraining on muscle strength and muscle power in multiple sclerosis patients. *NeuroRehabilitation* 2014; 34: 523–30.

9. Carter A, Daley A, Humphreys L, Snowden N, Woodroffe N, Petty J, et al. Pragmatic intervention for increasing self-directed exercise behaviour and improving important health outcomes in people with multiple sclerosis: a randomised controlled trial. *Mult Scler* 2014; 20: 1112–22.
10. Skjærbaek AG, Naesby M, Lutzen K, Møller AB, Jensen E, Lamers I, et al. Endurance training is feasible in severely disabled patients with progressive multiple sclerosis. *Mult Scler* 2014; 20: 627–30.
11. Briken S, Gold SM, Patra S, Vettorazzi E, Harbs D, Tallner A, et al. Effects of exercise on fitness and cognition in progressive MS: a randomized, controlled pilot trial. *Mult Scler* 2014; 20: 382–90.
12. Samaei A, Bakhtiari AH, Hajihasani A, Fatemi E, Motaharinezhad F. Uphill and Downhill Walking in Multiple Sclerosis: A Randomized Controlled Trial. *International journal of MS care* 2016; 18: 34–41.
13. Bansi J, Bloch W, Gamper U, Kesselring J. Training in MS: influence of two different endurance training protocols (aquatic versus overland) on cytokine and neurotrophin concentrations during three week randomized controlled trial. *Mult Scler* 2013; 19: 613–21.
14. Kargarfard M, Etemadifar M, Baker P, Mehrabi M, Hayatbakhsh R. Effect of aquatic exercise training on fatigue and health-related quality of life in patients with multiple sclerosis. *Arch Phys Med Rehabil* 2012; 93: 1701–8.
15. Wens I, Dalgas U, Vandenabeele F, Grevendonk L, Verboven K, Hansen D, et al. High Intensity Exercise in Multiple Sclerosis: Effects on Muscle Contractile Characteristics and Exercise Capacity, a Randomised Controlled Trial. *PloS one* 2015; 10: e0133697.
16. Hebert JR, Corboy JR, Manago MM, Schenkman M. Effects of vestibular rehabilitation on multiple sclerosis-related fatigue and upright postural control: a randomized controlled trial. *Phys Ther* 2011; 91: 1166–83.
17. Cattaneo D, Jonsdottir J, Zocchi M, Regola A. Effects of balance exercises on people with multiple sclerosis: a pilot study. *Clin Rehabil* 2007; 21: 771–81.
18. Kakt BD, Nacir B, Genc H, Saracoglu M, Karagoz A, Erdem HR, et al. Cycling progressive resistance training for people with multiple sclerosis: a randomized controlled study. *American journal of physical medicine & rehabilitation* 2010; 89: 446–57.
19. DeBolt LS, McCubbin JA. The effects of home-based resistance exercise on balance, power, and mobility in adults with multiple sclerosis. *Arch Phys Med Rehabil* 2004; 85: 290–7.
20. Forsberg A, von Koch L, Nilsagard Y. Effects on Balance and Walking with the CoDuSe Balance Exercise Program in People with Multiple Sclerosis: A Multicenter Randomized Controlled Trial. *Multiple sclerosis international* 2016; 2016: 7076265.
21. Hoang P, Schoene D, Gandevia S, Smith S, Lord SR. Effects of a home-based step training programme on balance, stepping, cognition and functional performance in people with multiple sclerosis—a randomized controlled trial. *Mult Scler* 2016; 22: 94–103.
22. Gandolfi M, Munari D, Geroi C, Gajofatto A, Benedetti MD, Midiri A, et al. Sensory integration balance training in patients with multiple sclerosis: A randomized, controlled trial. *Mult Scler* 2015; 21: 1453–62.
23. Bricchetto G, Piccardo E, Pedulla L, Battaglia MA, Tacchino A. Tailored balance exercises on people with multiple sclerosis: A pilot randomized, controlled study. *Mult Scler* 2015; 21: 1055–63.
24. Hilgers C, Mundermann A, Riehle H, Dettmers C. Effects of whole-body vibration training on physical function in patients with multiple sclerosis. *NeuroRehabilitation* 2013; 32: 655–63.
25. Kalron A, Rosenblum U, Frid L, Achiron A. Pilates exercise training vs. physical therapy for improving walking and balance in people with multiple sclerosis: a randomized controlled trial. *Clin Rehabil* 2017; 31: 319–28.
26. Fox EE, Hough AD, Creanor S, Gear M, Freeman JA. Effects of Pilates-Based Core Stability Training in Ambulant People With Multiple Sclerosis: Multicenter, Assessor-Blinded, Randomized Controlled Trial. *Phys Ther* 2016; 96: 1170–8.
27. Cramer H, Lauche R, Azizi H, Dobos G, Langhorst J. Yoga for multiple sclerosis: a systematic review and meta-analysis. *PloS one* 2014; 9: e112414.
28. Robinson J, Dixon J, Macsween A, van Schaik P, Martin D. The effects of exergaming on balance, gait, technology acceptance and flow experience in people with multiple sclerosis: a randomized controlled trial. *BMC sports science, medicine and rehabilitation* 2015; 7: 8.
29. Prosperini L, Fortuna D, Gianni C, Leonardi L, Marchetti MR, Pozzilli C. Home-based balance training using the Wii balance board: a randomized, crossover pilot study in multiple sclerosis. *NeuroRehabilitation and neural repair* 2013; 27: 516–25.
30. Nilsagard YE, Forsberg AS, von Koch L. Balance exercise for persons with multiple sclerosis using Wii games: a randomised, controlled multi-centre study. *Multiple sclerosis* *Mult Scler* 2013; 19: 209–16.
31. Pilutti LA, Dlugonski D, Sandroff BM, Klaren R, Motl RW. Randomized controlled trial of a behavioral intervention targeting symptoms and physical activity in multiple sclerosis. *Mult Scler* 2014; 20: 594–601.
32. Tarakci E, Yeldan I, Huseyinsinoglu BE, Zenginler Y, Eraksoy M. Group exercise training for balance, functional status, spasticity, fatigue and quality of life in multiple sclerosis: a randomized controlled trial. *Clin Rehabil* 2013; 27: 813–22.
33. Garrett M, Hogan N, Larkin A, Saunders J, Jakeman P, Coote S. Exercise in the community for people with minimal gait impairment due to MS: an assessor-blind randomized controlled trial. *Mult Scler* 2013; 19: 782–9.

34. Sosnoff JJ, Finlayson M, McAuley E, Morrison S, Motl RW. Home-based exercise program and fall-risk reduction in older adults with multiple sclerosis: phase 1 randomized controlled trial. *Clin Rehabil* 2014; 28: 254–63.
35. Heine M, van de Port I, Rietberg MB, van Wegen EE, Kwakkel G. Exercise therapy for fatigue in multiple sclerosis. *The Cochrane database of systematic reviews* 2015: Cd009956.
36. Dalgas U, Stenager E, Sloth M, Stenager E. The effect of exercise on depressive symptoms in multiple sclerosis based on a meta-analysis and critical review of the literature. *European journal of neurology* 2015; 22: 443–e34.
37. Broekmans T, Roelants M, Feys P, Alders G, Gijbels D, Hanssen I et al. Effects of long-term resistance training and simultaneous electro-stimulation on muscle strength and functional mobility in multiple sclerosis. *Mult Scler* 2011; 17: 468–77.
38. Dalgas U, Stenager E, Jakobsen J, Petersen T, Hansen HJ, Knudsen C et al. Resistance training improves muscle strength and functional capacity in multiple sclerosis. *Neurology* 2009; 73: 1478–84.
39. Dalgas U, Stenager E, Jakobsen J, Petersen T, Overgaard K, Ingemann-Hansen T. Muscle fiber size increases following resistance training in multiple sclerosis. *Mult Scler* 2010; 16: 1367–76.
40. Romberg A, Virtanen A, Ruutiainen J, Aunola S, Karppi SL, Vaara M et al. Effects of a 6-month exercise program on patients with multiple sclerosis: a randomized study. *Neurology* 2004; 63: 2034–8.
41. Surakka J, Romberg A, Ruutiainen J, Aunola S, Virtanen A, Karppi SL et al. Effects of aerobic and strength exercise on motor fatigue in men and women with multiple sclerosis: a randomized controlled trial. *Clin Rehabil* 2004; 18: 737–46.
42. Mostert S, Kesselring J. Effects of a short-term exercise training program on aerobic fitness, fatigue, health perception and activity level of subjects with multiple sclerosis. *Mult Scler* 2002; 8: 161–8.
43. Fimland MS, Helgerud J, Gruber M, Leivseth G, Hoff J. Enhanced neural drive after maximal strength training in multiple sclerosis patients. *Eur J Appl Physiol* 2010; 110: 435–43.
44. Beer S, Aschbacher B, Manoglou D, Gamper E, Kool J, Kesselring J. Robot-assisted gait training in multiple sclerosis: a pilot randomized trial. *Mult Scler* 2008; 14: 231–6.
45. Conklyn D, Stough D, Novak E, Paczak S, Chemali K, Bethoux F. A home-based walking program using rhythmic auditory stimulation improves gait performance in patients with multiple sclerosis: a pilot study. *NeuroRehabilitation and neural repair* 2010; 24: 835–42.
46. Snook EM, Motl RW. Effect of exercise training on walking mobility in multiple sclerosis: a meta-analysis. *NeuroRehabilitation and neural repair* 2009; 23: 108–16.
47. Pilutti LA, Dlugonski D, Sandroff BM, Klaren RE, Motl RW. Internet-delivered lifestyle physical activity intervention improves body composition in multiple sclerosis: preliminary evidence from a randomized controlled trial. *Arch Phys Med Rehabil* 2014; 95: 1283–8.
48. Borg GA. Psychophysical bases of perceived exertion. *Med Sci Sports Exerc* 1982; 14: 377–81.

Literatur zu Kapitel 37 Stürze

1. Schleicher MM, Wedam L, Wu G. Review of tai Chi as an effective exercise on falls prevention in elderly. *Res Sports Med* 2012; 20: 37–58.
2. Carter ND, Kannus P, Khan KM. Exercise in the prevention of falls in older people. A systematic literature review examining the rationale and the evidence. *Sports Med* 2001; 31: 427–38.
3. Gschwind YJ, Kressig RW, Lacroix A, Muehlbauer T, Pfenninger B, Granacher U. A best practice fall prevention exercise program to improve balance, strength/power, and psychosocial health in older adults: study protocol for a randomized controlled trial. *BMC Geriatr* 2013; 13: 105.
4. Shier V, Trieu E, Ganz DA. Implementing exercise programs to prevent falls: systematic descriptive review. *Inj Epidemiol* 2016; 3: 16.
5. Benichou O, Lord SR. Rationale for strengthening muscle to prevent falls and fractures: A review of the evidence. *Calcif Tissue Int* 2016; 98: 531–45.
6. Martin JT, Wolf A, Moore JL, Rolenz E, DiNinno A, Reneker JC. The effectiveness of physical therapist-administered group-based exercise on fall prevention: a systematic review of randomized controlled trials. *J Geriatr Phys Ther* 2013; 36: 182–93.
7. El-Khoury F, Cassou B, Charles MA, Dargent-Molina P. The effect of fall prevention exercise programmes on fall induced injuries in community dwelling older adults: systematic review and meta-analysis of randomised controlled trials. *BMJ* 2013; 347: f6234.
8. Kendrick D, Kumar A, Carpenter H, Zijlstra GA, Skelton DA, Cook JR et al. Exercise for reducing fear of falling in older people living in the community. *Cochrane Database Syst Rev* 2014; CD009848.
9. Reimers CD, Reimers N, Knapp G. Stürze. In: Reimers CD, Reuter I, Tettenborn B, Mewes N, Knapp G (Hrsg.) *Prävention und Therapie durch Sport. Band 2: Neurologie, Psychiatrie/Psychosomatik, Schmerzsyndrome. 2. Aufl., München: Elsevier Urban & Fischer, 2015, S. 143–161.*

Literatur zu Kapitel 38 Sarkopenie

1. Ali S, Garcia JM. Sarcopenia, cachexia and aging: diagnosis, mechanisms and therapeutic options – a mini-review. *Gerontology* 2014; 60: 294–305.
2. Csapo R, Alegre LM. Effects of resistance training with moderate vs heavy loads on muscle mass and strength in the elderly: A meta-analysis. *Scand J Med Sci Sports* 2016 Sep; 26: 995–1006. doi: 10.1111/sms.12536. Epub 2015 Aug 24.
3. Dodds R, Sayer AA. Sarcopenia. *Arq Bras Endocrinol Metabol* 2014 Jul; 58: 464–9.
4. Goisser S, Kemmler W, Porzel S, Volkert D, Sieber CC, Bollheimer LC, Freiburger E. Sarcopenic obesity and complex interventions with nutrition and exercise in community-dwelling older persons – a narrative review. *Clin Interv Aging* 2015 Aug 6; 10: 1267–82. doi: 10.2147/CIA.S82454. eCollection 2015.
5. Lee DC, Shook RP, Drenowatz C, Blair SN. Physical activity and sarcopenic obesity: definition, assessment, prevalence and mechanism. *Future Sci OA* 2016 Jul 14; 2: FSO127.
6. JafariNasabian P, Inglis JE, Kelly OJ, Ilich JZ. Osteosarcopenic obesity in women: impact, prevalence, and management challenges. *Int J Womens Health* 2017 Jan 13; 9: 33–42.
7. Law TD, Clark LA, Clark BC. Resistance Exercise to Prevent and Manage Sarcopenia and Dynapenia. *Annu Rev Gerontol Geriatr* 2016; 36: 205–228.
8. Morley JE. Frailty and Sarcopenia: The New Geriatric Giants. *Rev Invest Clin* 2016 Mar-Apr; 68: 59–67.
9. Padilla Colon CJ, Sanchez Collado P, Cuevas MJ. Beneficios del entrenamiento de fuerza para la prevención y tratamiento de la sarcopenia. *Nutr Hosp* 2014 May 1; 29: 979–88.
10. Reimers CD, Knapp G (2015) Sarkopenie. In: Reimers CD, Reuter I, Tettenborn B, Mewes N, Knapp G (Hrsg.) *Prävention und Therapie durch Sport. Band 2: Neurologie, Psychiatrie/Psychosomatik, Schmerzsyndrome*. 2. Aufl., Elsevier Urban & Fischer, München, S. 163–210.
11. Steffl M, Bohannon RW, Petr M, Kohlikova E, Holmerova I. Relation between cigarette smoking and sarcopenia: meta-analysis. *Physiol Res* 2015; 64: 419–26. Epub 2014 Dec 22.
12. Thomas DK, Quinn MA, Saunders DH, Greig CA. Protein Supplementation Does Not Significantly Augment the Effects of Resistance Exercise Training in Older Adults: A Systematic Review. *J Am Med Dir Assoc* 2016 Oct 1; 17: 959.e1–9. doi: 10.1016/j.jamda.2016.07.002.
13. Yu SC, Khaw KS, Jadcak AD, Visvanathan R. Clinical Screening Tools for Sarcopenia and Its Management. *Curr Gerontol Geriatr Res* 2016; 2016: 5978523. doi: 10.1155/2016/5978523. Epub 2016 Feb 4.
14. Zembron-Łacny A, Dziubek W, Rogowski Ł, Skorupka E, Dąbrowska G. Sarcopenia: monitoring, molecular mechanisms, and physical intervention. *Physiol Res* 2014; 63: 683–91.

Literatur zu Kapitel 39 Primäre Neuropathien und Polyneuropathien

1. Engelhardt A. Diagnostische Schritte bei Polyneuropathien. *Fortschr Neurol Psychiatr* 2008; 76: 429–439.
2. Heuß D, Auer-Grumbach M, Haupt FW, Löscher W, Neundörfer B, Rautenstraub B et al. Diagnostik bei Polyneuropathien. In: *Leitlinien für Diagnostik und Therapie in der Neurologie*. Stuttgart: Georg Thieme Verlag; 2008, S. 654 ff.
3. Ahn S, Song R. Effects of tai chi exercise on glucose control, neuropathy scores, balance, and quality of life in patients with type 2 diabetes and neuropathy. *J Altern Complement Med* 2012; 18: 1172–8.
4. Balducci S, Iacobellis G, Parisi L, Di Biase N, Calandriello E, Leonetti F et al. Exercise training can modify the natural history of diabetic peripheral neuropathy. *J Diabetes Complications* 2006; 20: 216–23.
5. Gong Q, Gregg EW, Wang J, An Y, Zhang P, Yang W et al. Long-term effects of a randomised trial of a 6-year lifestyle intervention in impaired glucose tolerance on diabetes-related microvascular complications: the China Da Qing Diabetes Prevention Outcome Study. *Diabetologia* 2011; 54: 300–7.
6. Smith AG, Russell J, Feldman EL, Goldstein J, Peltier A, Smith S et al. Lifestyle intervention for pre-diabetic neuropathy. *Diabetes Care* 2006; 29: 1294–9.
7. Dixit S, Maiya A, Shastri B. Effect of aerobic exercise on quality of life in population with diabetic peripheral neuropathy in type 2 diabetes: a single blind, randomized controlled trial. *Qual Life Res* 2013; 10.1007/s11136-013-0602-7.
8. Dixit S, Maiya AG, Shastri BA. Effect of aerobic exercise on peripheral nerve functions of population with diabetic peripheral neuropathy in type 2 diabetes: A single blind, parallel group randomized controlled trial. *J Diabetes And Its Complications* 2014; <http://dx.doi.org/10.1016/j.jdiacomp.2013.12.006>.
9. Mueller MJ, Tuttle LJ, LeMaster JW, Strube MJ, McGill JB, Hastings MK. Weight-bearing versus nonweight-bearing exercise for persons with diabetes and peripheral neuropathy: a randomized controlled trial. *Arch Phys Med Rehabil* 2013; 94: 829–38.
10. Grewal GS, Schwenk M, Lee-Eng J, Parvaneh S, Bharara M, Menzies RA, Talal TK, Armstrong DG, Najafi B. Sensor-Based Interactive Balance Training with Visual Joint Movement Feedback for Improving Postural Stability in Diabetics with

- Peripheral Neuropathy: A Randomized Controlled Trial. *Gerontology* 2015; 61: 567–74. doi: 10.1159/000371846. Epub 2015 Feb 19.
11. Chetlin RD, Gutmann L, Tarnopolsky M, Ullrich IH, Yeater RA. Resistance training effectiveness in patients with Charcot-Marie-Tooth disease: recommendations for exercise prescription. *Arch Phys Med Rehabil* 2004 Aug; 85: 1217–23.
 12. Chetlin RD, Gutmann L, Tarnopolsky MA, Ullrich IH, Yeater RA. Resistance training exercise and creatine in patients with Charcot-Marie-Tooth disease. *Muscle Nerve* 2004 Jul; 30: 69–76.
 13. Lindeman E, Leffers P, Reulen J, Spaans F, Drukker J. Quadriceps strength and timed motor performances in myotonic dystrophy, Charcot-Marie-Tooth disease, and healthy subjects. *Clin Rehabil* 1998 Apr; 12: 127–35.
 14. Lindeman E, Leffers P, Spaans F, Drukker J, Reulen J, Kerckhoffs M, Köke A. Strength training in patients with myotonic dystrophy and hereditary motor and sensory neuropathy: a randomized clinical trial. *Arch Phys Med Rehabil* 1995 Jul; 76: 612–20.
 15. Ramdharry GM, Pollard A, Anderson C, Laurá M, Murphy SM, Dudziec M, Dewar EL, Hutton E, Grant R, Reilly MM. A pilot study of proximal strength training in Charcot-Marie-Tooth disease. *J Peripher Nerv Syst* 2014 Dec; 19: 328–32. doi: 10.1111/jns.12100.
 16. Maggi G, Monti Bragadin M, Padua L, Fiorina E, Bellone E, Grandis M et al. Outcome measures and rehabilitation treatment in patients affected by Charcot-Marie-Tooth neuropathy: a pilot study. *Am J Phys Med Rehabil* 2011 Aug; 90(8): 628–37. doi: 10.1097/PHM.0b013e31821f6e32.
 17. El Mhandi L, Pichot V, Calmels P, Gautheron V, Roche F, Féasson L. Exercise training improves autonomic profiles in patients with Charcot-Marie-Tooth disease. *Muscle Nerve* 2011 Nov; 44: 732–6. doi: 10.1002/mus.22150.
 18. El Mhandi L, Millet GY, Calmels P, Richard A, Oullion R, Gautheron V, Féasson L. Benefits of interval-training on fatigue and functional capacities in Charcot-Marie-Tooth disease. *Muscle Nerve* 2008 May; 37: 601–10. doi: 10.1002/mus.20959.
 19. Mkandla K, Myezwa H, Musenge E. The effects of progressive-resisted exercises on muscle strength and health-related quality of life in persons with HIV-related poly-neuropathy in Zimbabwe. *AIDS Care* 2016 May; 28: 639–43. doi: 10.1080/09540121.2015.1125418. Epub 2016 Jan 5.
 20. Schwenk M, Grewal GS, Holloway D, Muchna A, Garland L, Najafi B. Interactive Sensor-Based Balance Training in Older Cancer Patients with Chemotherapy-Induced Peripheral Neuropathy: A Randomized Controlled Trial. *Gerontology* 2015 Dec 18 (Epub ahead of print).

Literatur zu Kapitel 40 Spinale Muskelatrophien

1. Montes J, Garber CE, Kramer SS, Montgomery MJ, Dunaway S et al. Single-Blind, Randomized, Controlled Clinical Trial of Exercise in Ambulatory Spinal Muscular Atrophy: Why are the Results Negative? *J Neuromuscul Dis* 2015; 2: 463–70.
2. Lewelt A, Krossschell KJ, Stoddard GJ, Weng C, Xue M, et al. Resistance strength training exercise in children with spinal muscular atrophy. *Muscle Nerve* 2015; 52: 559–67.
3. Shrader JA, Kats I, Kokkinis A, Zampieri C, Levy E, et al. A randomized controlled trial of exercise in spinal and bulbar muscular atrophy. *Ann Clin Transl Neurol* 2015; 2:739–47.
4. Dahlqvist JR and Vissing J. Exercise Therapy in Spinobulbar Muscular Atrophy and Other Neuromuscular Disorders. *J Mol.Neurosci* 2016; 58: 388–93.

Literatur zu Kapitel 41 Postpoliomyelitis-Syndrom

1. Koopman FS, Voorn EL, Beelen A, Bleijenberg G, de Visser M, et al. No Reduction of Severe Fatigue in Patients With Postpolio Syndrome by Exercise Therapy or Cognitive Behavioral Therapy: Results of an RCT. *Neurorehabil Neural Repair* 2016; 30: 402–10.

Literatur zu Kapitel 42 Myopathien

1. Schupp W. Hereditäre Myopathien. In: Reimers CD, Reuter I, Tettenborn B, Brooks A, Thürauf N, Knapp G CD (Hrsg.) *Prävention und Therapie durch Sport. Band 2. Neurologie, Psychiatrie/Psychosomatik, Schmerzsyndrome.* 2. Aufl., München: Urban & Fischer, 2016, S. 235–51.
2. Matthews E, Brassington R, Kuntzer T, Jichi F, Manzur AY. Corticosteroids for the treatment of Duchenne muscular dystrophy. *Cochrane Database Syst Rev.* 2016; CD003725.

3. Andersen G, Prahm KP, Dahlqvist JR, Citirak G, Vissing J: Aerobic training and postexercise protein in facioscapulohumeral muscular dystrophy. RCT study. *Neurology*. 2015; 85: 396–403.
4. Siciliano G, Simoncini C, Giannotti S, Zampa V, Angelini C, Ricci G. Muscle exercise in limb girdle muscular dystrophies: pitfall and advantages. *Acta Myol*. 2015; 34: 3–8.
5. Vissing CR, Preisler N, Prahm KP, Vissing J: aerobic training in patients with anoctamin 5 myopathy and hyperckemia. *Musclr Nerve*. 2014; 50: 119–23.
6. Brady LI, MacNeil JG, Tarnopolsky MA: Impact of habitual exercise on the strength of individuals with myotonic dystrophy type 1. *Am J Phys Med Rehabil*. 2014; 93: 739–50.
7. Hedermann G, Vissing CR, Heje K, Preisler N, Witting N, Vissing J. Aerobic training in patients with congenital myopathy. *PLoS One*. 2016; 11: e0146036.
8. Preisler N, Haller RG, Vissing J: Exercise in muscle glycogen storage diseases. *J Inherit Metab Dis*. 2015;38:551–63.
9. Tarnopolsky MA: Exercise as a therapeutic strategy for primary mitochondrial strategies. *J Child Neurol*. 2014; 29: 1225–34.
10. Kwak HB. Statin-induced myopathy in skeletal muscle: the role of exercise. *J Lifestyle Med*. 2014; 4: 71–9.

Literatur zu Kapitel 43 Entzündliche Muskelerkrankungen

1. Dalakas MC, Hohlfeld R. Polymyositis and dermatomyositis. *Lancet* 2003; 362: 971–82.
2. Albayda J, Christopher-Stine L. Novel approaches in the treatment of myositis and myopathies. *Ther Adv Musculoskelet Dis* 2012; 4: 369–77.
3. Cherin P. Recognition and management of myositis. *Drugs* 1997; 54: 39–49.
4. Griggs RC. The current status of treatment for inclusion-body myositis. *Neurology* 2006; 66 (2 Suppl 1): S30–2.
5. Walter MC et al. High-dose immunoglobulin therapy in sporadic inclusion body myositis: a double-blind, placebo-controlled study. *J Neurol* 2000; 247: 22–8.
6. Marie I. et al. Polymyositis and dermatomyositis: short term and longterm outcome, and predictive factors of prognosis. *J Rheumatol* 2001; 28: 2230–7.
7. Wiesinger GF et al. Aerobic capacity in adult dermatomyositis/polymyositis patients and healthy controls. *Arch Phys Med Rehabil* 2000; 81: 1–5.
8. Chung YL et al. Creatine supplements in patients with idiopathic inflammatory myopathies who are clinically weak after conventional pharmacologic treatment: Six-month, double-blind, randomized, placebo-controlled trial. *Arthritis Rheum* 2007; 57: 694–702.
9. Wiesinger GF et al. Benefit of 6 months long-term physical training in polymyositis/dermatomyositis patients. *Br J Rheumatol* 1998; 37: 1338–42.
10. Mathiesen PR et al. Aerobic fitness after JDM—a long-term follow-up study. *Rheumatology* 2013; 52: 287–95.
11. Arnardottir S et al. Sporadic inclusion body myositis: pilot study on the effects of a home exercise program on muscle function, histopathology and inflammatory reaction. *J Rehabil Med* 2003; 35: 31–5.
12. Escalante A, Miller L, Beardmore TD. Resistive exercise in the rehabilitation of polymyositis/dermatomyositis. *J Rheumatol* 1993; 20: 1340–4.
13. Keteyian SJ. Cardiopulmonary adaption to exercise. *ACSM's Resource Manual for Guidelines for Exercise Testing and Prescription*. ed. L.A. Kaminsky. Vol. 5th edition. 2006, Philadelphia, PA: Lippincott Williams and Wilkins. 13.
14. Alemo Munters L et al. Improved exercise performance and increased aerobic capacity after endurance training of patients with stable polymyositis and dermatomyositis. *Arthritis Res Ther* 2013; 15: R83.
15. Nader GA et al. A longitudinal, integrated, clinical, histological and mRNA profiling study of resistance exercise in myositis. *Mol Med* 2010; 16: 455–64.
16. Spector SA et al. Safety and efficacy of strength training in patients with sporadic inclusion body myositis. *Muscle Nerve* 1997; 20: 1242–8.

Literatur zu Kapitel 44 Myasthenia gravis

1. Lindstrom J. An assay for antibodies to human acetylcholine receptor in serum from patients with myasthenia gravis. *Clin Immunol Immunopathol* 1977; 7: 36–43.
2. Vincent A, Newsom-Davis J. Acetylcholine receptor antibody as a diagnostic test for myasthenia gravis: results in 153 validated cases and 2967 diagnostic assays. *J Neurol Neurosurg Psychiatry* 1985; 48: 1246–52.

3. Evoli A et al. Clinical correlates with anti-MuSK antibodies in generalized seronegative myasthenia gravis. *Brain* 2003; 126: 2304–11.
4. McConville J et al. Detection and characterization of MuSK antibodies in seronegative myasthenia gravis. *Ann Neurol* 2004; 55: 580–4.
5. Somnier FE, Engel PJ. The occurrence of anti-titin antibodies and thymomas: a population survey of MG 1970–1999. *Neurology* 2002; 59: 92–8.
6. Yamamoto AM et al. Anti-titin antibodies in myasthenia gravis: tight association with thymoma and heterogeneity of nonthymoma patients. *Arch Neurol* 2001; 58: 885–90.
7. Cavalcante P, Bernasconi P, Mantegazza R. Autoimmune mechanisms in myasthenia gravis. *Curr Opin Neurol* 2012; 25: 621–9.
8. Namba T, Brunner NG, Grob D. Myasthenia gravis in patients with thymoma, with particular reference to onset after thymectomy. *Medicine* 1978; 57: 411–33.
9. Juel VC, Massey JM. Myasthenia gravis. *Orphanet J Rare Dis* 2007; 2: 44.
10. Fregonezi GA et al. Effects of 8-week, interval-based inspiratory muscle training and breathing retraining in patients with generalized myasthenia gravis. *Chest* 2005; 128: 1524–30.
11. Phillips LH. The epidemiology of myasthenia gravis. *Ann N Y Acad Sci* 2003; 998: 407–12.
12. Sanders DB et al. Clinical aspects of MuSK antibody positive seronegative MG. *Neurology* 2003; 60: 1978–80.
13. Pardy RL, Hussain SN, Macklem PT. The ventilatory pump in exercise. *Clin Chest Med* 1984; 5: 35–49.
14. Shaffer TH, Wolfson MR, Bhutani VK. Respiratory muscle function, assessment, and training. *Phys Ther* 1981; 61: 1711–23.
15. Leith DE, Bradley M. Ventilatory muscle strength and endurance training. *J Appl Physiol* 1976; 41: 508–16.
16. Keens TG et al. Ventilatory muscle endurance training in normal subjects and patients with cystic fibrosis. *Am Rev Respir Dis* 1977; 116: 853–60.
17. Weiner P et al. Inspiratory muscle training in patients with bronchial asthma. *Chest* 1992; 102: 1357–61.
18. Pardy RL et al. Inspiratory muscle training compared with physiotherapy in patients with chronic airflow limitation. *Am Rev Respir Dis* 1981; 123: 421–5.
19. Hernandez MT et al. Results of a home-based training program for patients with COPD. *Chest* 2000; 118: 106–14.
20. Lotters F et al. Effects of controlled inspiratory muscle training in patients with COPD: a meta-analysis. *Eur Respir J* 2002; 20: 570–6.
21. Klefbeck B, Lagerstrand L, Mattsson E. Inspiratory muscle training in patients with prior polio who use part-time assisted ventilation. *Arch Phys Med Rehabil* 2000; 81: 1065–71.
22. Gross D et al. The effect of training on strength and endurance of the diaphragm in quadriplegia. *Am J Med* 1980; 68: 27–35.
23. Liaw MY et al. Resistive inspiratory muscle training: its effectiveness in patients with acute complete cervical cord injury. *Arch Phys Med Rehabil* 2000; 81: 752–6.
24. Rassler B et al. The effect of respiratory muscle endurance training in patients with myasthenia gravis. *Neuromuscul Disord* 2007; 17: 385–91.
25. Weiner P et al. Respiratory muscle training in patients with moderate to severe myasthenia gravis. *Can J Neurol Sci* 1998; 25: 236–41.
26. Lohi EL, Lindberg C, Andersen O. Physical training effects in myasthenia gravis. *Arch Phys Med Rehabil* 1993; 74: 1178–80.
27. Stout JR et al. Effects of resistance exercise and creatine supplementation on myasthenia gravis: a case study. *Med Sci Sports Exerc* 2001; 33: 869–72.

Literatur zu Kapitel 45 Kognitive Störungen

1. Felbecker A, Tettenborn B, Reimers CD, Knapp G. Kognitive Störungen. In: *Prävention und Therapie durch Sport*. Band 2: Neurologie, Psychiatrie/Psychosomatik, Schmerzsyndrome. 2. Auflage 2015, Elsevier GmbH München.
2. Wu YT, Fratiglioni L, Matthews FE, Lobo A, Breteler MM, et al. Dementia in western Europe: epidemiological evidence and implications for policy making. *Lancet Neurol* 2016; 15: 116–24.
3. Satizabal CL, Beiser AS, Chouraki V, Chêne G, Dufouil C, Seshadri S. Incidence of Dementia over Three Decades in the Framingham Heart Study. *N Engl J Med* 2016; 374: 523–32.
4. Felbecker A, Tettenborn B. Rolle vaskulärer Risikofaktoren für die Entstehung und Progression der Alzheimer-Demenz. *Akt Neurol* 2016; 43: 309–17.
5. Moll van Charante EP, Richard E, Eurelings LS, van Dalen J-W, Ligthart SA et al. Effectiveness of a 6-year multidomain vascular care intervention to prevent dementia (preDIVA): a cluster-randomised controlled trial. *Lancet* 2016; 388: 797–805.

6. Ngandu T, Lehtisalo J, Solomon A, Levälähti E, Ahtiluoto S, et al. A 2 year multidomain intervention of diet, exercise, cognitive training, and vascular risk monitoring versus control to prevent cognitive decline in at-risk elderly people (FINGER): a randomised controlled trial. *Lancet* 2015; 385: 2255–63.
7. Wagner G, Herbsleb M, Cruz F, Schumann A, Köhler S, et al. Changes in fMRI activation in anterior hippocampus and motor cortex during memory retrieval after an intense exercise intervention. *Biol Psychol* 2017; doi: 10.1016/j.biopsycho.2017.01.003.
8. Sabia S, Dugravot A, Dartigues JF, Abell J, Elbaz A, Kivimäki M, Singh-Manoux A. Physical activity, cognitive decline, and risk of dementia: 28 year follow-up of Whitehall II cohort study. *BMJ* 2017 Jun 22; 357: j2709. doi: 10.1136/bmj.j2709
9. Barnes DE, Yaffe K. The projected effect of risk factor reduction on Alzheimer's disease prevalence. *Lancet Neurol* 2011; 10: 819–28.
10. Groot C, Hooghiemstra AM, Raijmakers PG, van Berckel BN, Scheltens P, et al. The effect of physical activity on cognitive function in patients with dementia: A meta-analysis of randomized control trials. *Ageing Res Rev* 2016; 25: 13–23.
11. Train the Brain Consortium. Randomized trial on the effects of a combined physical/cognitive training in aged MCI subjects: the Train the Brain study. *Nature Sci Rep* 2017; 7: 39471.
12. Stephen R, Hongisto K, Solomon A, Lönnroos E. Physical activity and Alzheimer's disease: a systematic review. *J Gerontol A Biol Sci Med Sci* 2017; 3: 733–739.
13. Henwood T, Neville C, Baguley C, Beattie E. Aquatic exercise for residential aged care adults with dementia: benefits and barriers to participation. *Int Psychogeriatr* 2017; 29: 1439–1449.

Literatur zu Kapitel 46 Depression

1. Brooks A, Pink J. Voraussetzungen für eine erfolgreiche Pharmakotherapie von Depressionen. *DNP – der Neurologe und Psychiater* 2016; 17: 7–8.
2. Schuch FB, Vancampfort D, Sui X, Rosenbaum S, Firth J, Richards J, Ward PB, Stubbs B. Are lower levels of cardiorespiratory fitness associated with incident depression? A systematic review of prospective cohort studies. *Prev Med* 2016 Dec; 93: 159–165.
3. Farmer ME, Locke BZ, Mościcki EK, Dannenberg AL, Larson DB, Radloff LS. Physical activity and depressive symptoms: the NHANES I Epidemiologic Follow-up Study. *Am J Epidemiol* 1988 Dec; 128: 1340–51.
4. Greist JH, Klein MH, Eischens RR, Faris J, Gurman AS, Morgan WP. Running as treatment for depression. *Compr Psychiat* 1979; 20: 41–54.
5. Martinsen EW, Medhus A, Sandvik L. Effects of aerobic exercise on depression: a controlled study. *Brit Med J* 1985; 291: 109.
6. Martinsen EW, Hoffart A, Solberg O. Comparing aerobic with anaerobic forms of exercise in the treatment of clinical depression: a randomized trial. *Compr Psychiatry* 1989; 30: 324–31.
7. Blumenthal JA, Babyak MA, Moore KA, Craighead WE, Herman S et al. Effects of exercise training on older patients with major depression. *Arch Intern Med* 1999; 159: 2349–56.
8. Babyak M, Blumenthal J, Herman S. Exercise treatment for major depression: maintenance of therapeutic benefit over 10 months. *Psychosom Med* 2000; 62: 633–8.
9. Rimer J, Dwan K, Lawlor DA, Greig CA, McMurdo M, Morley W, Mead GE. Exercise for depression. *Cochrane Database Syst Rev* 2012; 7.
10. Rethorst CD, Wipfli BM, Landers DM. The antidepressive effects of exercise. A meta-analysis of randomized trials. *Sports Med* 2009; 39: 491–511.
11. Krogh J, Nordentoft M, Sterne JA, Lawlor DA. The effect of exercise in clinically depressed adults: systematic review and meta-analysis of randomized controlled trials. *J Clin Psychiatry* 2011; 72: 529–38.
12. Silveira H, Moraes H, Oliveira N, Coutinho ES, Laks J, Deslandes A. Physical exercise and clinically depressed patients: a systematic review and meta-analysis. *Neuropsychobiology* 2013; 67: 61–8.
13. Bridle C, Spanjers K, Patel S, Atherton NM, Lamb SE. Effect of exercise on depression severity in older people: systematic review and meta-analysis of randomised controlled trials. *Br J Psychiatry* 2012; 201: 180–5.
14. Meyer JD, Koltyn KF, Stegner AJ, Kim JS, Cook DB. Relationships between serum BDNF and the antidepressant effect of acute exercise in depressed women. *Psychoneuroendocrinology*. 2016 Sep 28; 74: 286–294.
15. Brooks A, Rieckmann P. Sport und Neuroplastizität bei psychischen Erkrankungen. *Info Neurologie und Psychiatrie* 2014; 16: 51–59.
16. Schuch FB, Vancampfort D, Richards J, Rosenbaum S, Ward PB, Stubbs B. Exercise as a treatment for depression: A meta-analysis adjusting for publication bias. *J Psychiatr Res* 2016 Jun; 77: 42–51.
17. Voderholzer U, Dersch R, Dickhuth HH, Herter A, Freyer T, Berger M. Physical fitness in depressive patients and impact of illness course and disability. *J Affect Disord* 2011; 128: 160–4.

18. Blumenthal JA, Babyak MA, Doraiswamy PM, Watkins L, Hoffman BM, Barbour KA, Herman S, Craighead WE, Brosse AL, Waugh R, Hinderliter A, Sherwood A. Exercise and pharmacotherapy in the treatment of major depressive disorder. *Psychosom Med* 2007; 69: 587–96.
19. Dunn AL, Trivedi MH, Kampert JB, Clark CG, Chambliss HO. Exercise treatment for depression: efficacy and dose response. *Am J Prev Med* 2005; 28: 1–8.
20. Dunn AL, Trivedi MH, Kampert JB, Clark CG, Chambliss HO. The DOSE study: a clinical trial to examine efficacy and dose response of exercise as treatment for depression. *Control Clin Trials* 2002; 23: 584–603.

Literatur zu Kapitel 47 Angststörungen

1. Berger M. Psychische Erkrankungen: Klinik und Therapie. München: Elsevier Urban & Fischer, 2014.
2. Broocks A. Körperliche Aktivität und psychische Gesundheit – Psychische und neurobiologische Effekte von Ausdauertraining bei Patienten mit Panikstörung und Agoraphobie. Monographien aus dem Gesamtgebiete der Psychiatrie. Darmstadt: Steinkopff, 2000.
3. Ströhle A, Feller C, Onken M, Godemann F, Heinz A, Dimeo F. The acute antipanic activity of aerobic exercise. *Am J Psychiatry* 2005; 162: 2376–8.
4. Ströhle A, Graetz B, Scheel M, Wittmann A, Feller C et al. The acute antipanic and anxiolytic activity of aerobic exercise in patients with panic disorder and healthy control subjects. *J Psychiatr Res* 2009; 43: 1013–7.
5. Broocks A, Bandelow B, Pekrun G, George A, Meyer T et al. Comparison of aerobic exercise, clomipramine, and placebo in the treatment of panic disorder. *Am J Psychiatry* 1998; 155: 603–9.
6. Wedekind D, Broocks A, Weiss N, Engel K, Neubert K, Bandelow B. A randomized, controlled trial of aerobic exercise in combination with paroxetine in the treatment of panic disorder. *World J Biol Psychiatry* 2010; 11: 904–13.
7. Herring MP, Jacob ML, Suveg C, Dishman RK, O'Connor PJ. Feasibility of exercise training for the short-term treatment of generalized anxiety disorder: a randomized controlled trial. *Psychother Psychosom* 2012; 81: 21–8.
8. Wipfli BM, Rethorst CD, Landers DM. The anxiolytic effects of exercise: a meta-analysis of randomized trials and dose-response analysis. *J Sport Exerc Psychol*. 2008 Aug; 30 : 392–410. Erratum in: *J Sport Exerc Psychol* 2009; 31: 128–9.
9. Jorm AF, Christensen H, Griffiths KM, Parslow RA, Rodgers B, Blewitt KA. Effectiveness of complementary and self-help treatments for anxiety disorders. *Med J Aust* 2004; 181 (7 Suppl): S29–46.
10. Brown RA, Abrantes AM, Strong DR, Mancebo MC, Menard J et al. A pilot study of moderate-intensity aerobic exercise for obsessive compulsive disorder. *J Nerv Ment Dis* 2007; 195: 514–20.
11. Abrantes AM, Strong DR, Cohn A, Cameron AY, Greenberg BD et al. Acute changes in obsessions and compulsions following moderate-intensity aerobic exercise among patients with obsessive-compulsive disorder. *J Anxiety Disord* 2009; 23: 923–7.
12. Abrantes AM, McLaughlin N, Greenberg BD, Strong DR, Riebe D et al. Design and rationale for a randomized controlled trial testing the efficacy of aerobic exercise for patients with obsessive-compulsive disorder. *Ment Health Phys Act* 2012; 5: 155–65.
13. Jayakody K, Gunadasa S, Hosker C. Exercise for anxiety disorders: systematic review. *Br J Sports Med* 2014 Feb; 48: 187–96.

Literatur zu Kapitel 48 Psychosen aus dem schizophrenen Formenkreis

1. Möller H-J, Laux G, Kapfhammer H-P (Hrsg). Lehrbuch Psychiatrie, Psychosomatik und Psychotherapie. 4. Aufl., Heidelberg: Springer 2010.
2. Vancampfort D, Probst M, Scheewe T, De Herdt A, Sweers K et al. Relationships between physical fitness, physical activity, smoking and metabolic and mental health parameters in people with schizophrenia. *Psychiatry Res* 2013; 207: 25–32.
3. Beebe LH, Smith K, Burk R, McIntyre K, Dessieux O et al. Effect of a motivational intervention on exercise behavior in persons with schizophrenia spectrum disorders. *Community Ment Health J* 2011; 47: 628–36.
4. Naber D, Moritz S, Lambert M, Pajonk FG, Holzbach R et al. Improvement of schizophrenic patients' subjective well-being under atypical antipsychotic drugs. *Schizophr Res* 2001; 50: 79–88.
5. Deimel H, Lohmann S. Physical capacity of schizophrenic patients. *Rehabilitation* 1983; 22: 81–5.
6. Vancampfort D, Probst M, De Herdt A, Corredeira RM, Carraro A et al. An impaired health related muscular fitness contributes to a reduced walking capacity in patients with schizophrenia: a cross-sectional study. *BMC Psychiatry* 2013; 13: 5.
7. Pajonk FG, Wobrock T, Gruber O, Scherk H, Berner D, Kaizl I et al. Hippocampal plasticity in response to exercise in schizophrenia. *Arch Gen Psychiatry* 2010; 67: 133–43.

8. Scheewe TW, Backx FJ, Takken T, Jörg F, van Strater AC et al. Exercise therapy improves mental and physical health in schizophrenia: a randomised controlled trial. *Acta Psychiatr Scand* 2013; 127: 464–73.
9. Scheewe TW, van Haren NE, Sarkisyan G, Schnack HG, Brouwer RM et al. Exercise therapy, cardiorespiratory fitness and their effect on brain volumes: A randomised controlled trial in patients with schizophrenia and healthy controls. *Eur Neuropsychopharmacol* 2012 Sep 12-
10. Firth J, Carney R, Elliott R, French P, Parker S, McIntyre R, McPhee JS, Yung AR. Exercise as an intervention for first-episode psychosis: a feasibility study. *Early Interv Psychiatry* 2016 Mar 14.
11. Firth J, Stubbs B, Rosenbaum S, Vancampfort D, Malchow B, Schuch F et al. Aerobic Exercise Improves Cognitive Functioning in People With Schizophrenia: A Systematic Review and Meta-Analysis. *Schizophr Bull* 2016 Aug 11. pii: sbw115. [Epub ahead of print].
12. Brönnner M, Betz C, Christ S, Froböse T, Pitschel-Walz G, Bäuml J. Wellness-activities to prevent metabolic syndrome among patients with schizophrenia. *Psychiatr Prax* 2010; 37: 7–19.
13. Daumit GL, Dickerson FB, Wang NY, Dalcin A, Jerome GJ et al. A behavioral weight-loss intervention in persons with serious mental illness. *N Engl J Med* 2013; 368: 1594–602.
14. Firth J, Rosenbaum S, Stubbs B, Gorczynski P, Yung AR, Vancampfort D. Motivating factors and barriers towards exercise in severe mental illness: a systematic review and meta-analysis. *Psychol Med* 2016 Oct; 46: 2869–2881. Epub 2016 Aug 9.
15. Browne J, Penn DL, Battaglini CL, Ludwig K. Work out by Walking: A Pilot Exercise Program for Individuals With Schizophrenia Spectrum Disorders. *J Nerv Ment Dis* 2016; 204: 651–7.

Literatur zu Kapitel 49 Abhängigkeitserkrankungen

1. Pabst A. et al. Substanzkonsum und substanzbezogene Störungen in Deutschland im Jahre 2012. *Sucht* 2013; 59: 321–331.
2. Schneider U, Altmann A, Baumann M, Bernzen J, Bertz B et al. Comorbid anxiety and affective disorder in alcohol-dependent patients seeking treatment: the first Multicentre Study in Germany. *Alcohol Alcohol* 2001; 36: 219–23.
3. Driessen M, Meier S, Hill A, Wetterling T, Lange W, Junghanns K. The course of anxiety, depression and drinking behaviours after completed detoxification in alcoholics with and without comorbid anxiety and depressive disorders. *Alcohol Alcohol* 2000; 36: 249–55.
4. Miller WR, Rollnick S. Motivierende Gesprächsführung. Ein Konzept zur Beratung von Menschen mit Suchtproblemen. Freiburg: Lambertus 1999.
5. Swift RM. Medications and alcohol craving. *Alcohol Res Health* 1999; 23: 207–13.
6. Feeney GF, Young RM, Connor JP, Tucker J, McPherson A. Outpatient cognitive behavioural therapy programme for alcohol dependence: impact of naltrexone use on outcome. *Aust N Z J Psychiatry* 2001; 35: 443–8.
7. Stuppe M, Froesa G, Pinzke R. Community Reinforcement Approach in einer psychiatrischen Klinik. *Suchttherapie* 2013; 14: 56–63.
8. Kirkcaldy BD, Shephard RJ, Siefen RG. The relationship between physical activity and self-image and problem behaviour among adolescents. *Soc Psychiatry Psychiatr Epidemiol* 2002; 37: 544–550.
9. Dakwar E, Blanco C, Lin KH, Liu SM, Warden D et al. Exercise and mental illness: results from the National Epidemiologic Survey on Alcohol and Related Conditions (NESARC). *J Clin Psychiatry* 2012; 73: 960–6.
10. Sinyor D, Brown T, Rostant L, Seraganian P. The role of a physical fitness program in the treatment of alcoholism. *J Studies Alcohol* 1982; 43: 380–6.
11. Bartmann U. Laufen und Joggen und seine positive Auswirkung auf die Psyche: Stuttgart: TRIAS – Thieme, Hippokrates, Enke 1991.
12. Brown RA, Abrantes AM, Minami H, Read JP, Marcus BH, Jakicic JM et al. A preliminary, randomized trial of aerobic exercise for alcohol dependence. *J Subst Abuse Treat* 2014 Jul; 47: 1–9.
13. Zschucke E, Heinz A, Ströhle A. Exercise and physical activity in the therapy of substance use disorders. *Scientific World Journal* 2012; 901:741. doi: 10.1100/2012/901741. Epub 2012 May 3. Review.
14. Giesen ES, Deimel H, Bloch W. Clinical exercise interventions in alcohol use disorders: a systematic review. *J Subst Abuse Treat* 2015 May; 52: 1–9.
15. Manthou E, Georgakouli K, Fatouros IG, Gianoulakis C, Theodorakis Y, Jamurtas AZ. Role of exercise in the treatment of alcohol use disorders. *Biomed Rep* 2016; 4: 535–545.
16. Breslau N, Kilbey M, Andreski P. Nicotine dependence, major depression and anxiety in young adults. *Arch Gen Psychiatry* 1991; 48: 1069–74.
17. Kröger CB, Lohmann B. Tabakkonsum und Tabakabhängigkeit. *Hogrefe* 2007, S. 122.
18. Glassman AH. Cigarette smoking: Implications for psychiatric illness. *Am J Psychiatry* 1993; 150: 546–53.

19. Mann-Wrobel MC, Bennett ME, Weiner EE, Buchanan RW, Ball MP. Smoking history and motivation to quit in smokers with schizophrenia in a smoking cessation program. *Schizophr Res* 2011 Mar; 126: 277–83.
20. Weinberger DR, Bergman KF, Illowski BP. Physiological dysfunction of dorsolateral prefrontal cortex in schizophrenia, III: a new cohort and evidence for a monoaminergic mechanism. *Arch Gen Psych* 1988; 45: 609–15.
21. Carr A. Endlich Nichtraucher! Der einfache Weg, mit dem Rauchen Schluss zu machen. München: Goldmann, 2012.
22. Lohmann B. Frau Schmitt, die Zigaretten und Sie: Ratgeber zur Tabakentwöhnung. Göttingen: Hogrefe 2009.
23. Batra A, Mann K. Alkohol und Tabak: Grundlagen und Folgeerkrankungen. Stuttgart: Thieme, 2010
24. Marcus BH, Albrecht AE, Niaura RS, Abrams DB, Thompson PD. Usefulness of physical exercise for maintaining smoking cessation in women. *Am J Cardiol* 1991; 68: 406–7.
25. Taylor CB, Houston Miller N, Haskell WL, Debusk RF. Smoking cessation after acute myocardial infarction: the effects of exercise training. *Addict Behav* 1988; 13: 331–5.
26. Haasova M, Warren FC, Ussher M, Janse Van Rensburg K et al. The acute effects of physical activity on cigarette cravings: systematic review and meta-analysis with individual participant data. *Addiction* 2013; 108: 26–37.
27. Ussher MH, Taylor A, Faulkner G. Exercise interventions for smoking cessation. *Cochrane Database Syst Rev* 2012; 18;1:CD002295.
28. Berger M. Psychische Erkrankungen: Klinik und Therapie. München: Elsevier Urban & Fischer, 2011.
29. Zschucke E, Heinz A, Ströhle A. Exercise and physical activity in the therapy of substance use disorders. *Scientific World Journal* 2012; 901: 741.
30. Wang D, Zhou C, Zhao M, Wu X, Chang YK. Dose-response relationships between exercise intensity, cravings, and inhibitory control in methamphetamine dependence: An ERPs study. *Drug Alcohol Depend* 2016 Apr 1; 161: 331–9.
31. Sabine M. Grüsser, Carolin N. Thalemann: Verhaltenssucht. Diagnostik, Therapie, Forschung. Bern: Huber, 2006.
32. Derbyshire KL, Lust KA, Schreiber LR, Odlaug BL, Christenson GA, Golden DJ, Grant JE. Problematic Internet use and associated risks in a college sample *Compr Psychiatry* 2013 Jul; 54: 415–22.
33. Glasser W. Positive Addiction. New York, Evanston, San Francisco, London: Harper & Row, 1976.
34. Fagerstrom K, Furberg H. A comparison of the Fagerström Test for Nicotine Dependence and smoking prevalence across countries. *Addiction* 2008; 103: 841–5.
35. Heinz A, Batra A. Neurobiologie der Alkohol- und Nikotinabhängigkeit. Stuttgart: Kohlhammer, 2003.
36. Lehner B, Kepp J. Daten, Zahlen, Fakten. In: Deutsche Hauptstelle für Suchtfragen: Jahrbuch Sucht. Pabst Lengerich 2016.

Literatur zu Kapitel 50 Kopfschmerzen

1. Classification Committee of the International Headache Society. The International Headache Disorders, 2nd ed. Cephalgia 2004; 24 (Suppl 1): 9–160.
2. Evers S, May A, Fritsche G, Kropp P, Lampl C, Limmroth V, Malzacher V, Sandor S, Straube A, Diener HC: Akuttherapie und Prophylaxe der Migräne – Leitlinie der Deutschen Migräne- und Kopfschmerzgesellschaft und der Deutschen Gesellschaft für Neurologie. *Nervenheilkunde* 2008; 27: 933–49.
3. Chen Y. Advances in the pathophysiology of tension-type headache: from stress to central sensitization. *Curr Pain Headache Rep* 2009; 13: 484–94.
4. Leitlinien DGN 2016. A. Straube. Therapie des episodischen und chronischen Kopfschmerz vom Spannungstyp und anderer chronischer täglicher Kopfschmerzen. AWMF-Registernummer 030/077.
5. Fernández-de-las-Peñas C, Alonso-Blanco C, San-Roman J, Miangolarra-Page JC. Methodological quality of randomized controlled trials of spinal manipulation and mobilization in tension-type headache, migraine, and cervicogenic headache. *Orthop Sports Phys Ther* 2006; 36: 160–9.
6. Castien RF, van der Windt DA, Grooten A, Dekker J. Effectiveness of manual therapy for chronic tension-type headache: a pragmatic, randomised, clinical trial. *Cephalalgia* 2011; 31: 133–43.
7. Torelli P, Jensen R, Olesen J. Physiotherapy for tension-type headache: a controlled study. *Cephalalgia* 2004; 24: 29–36.
8. Linde K, Allais G, Brinkhaus B, Manheimer E, Vickers A, White AR. Acupuncture for tension-type headache. *Cochrane Database Syst Rev* 2009; 21: CD007587.
9. Bendtsen L, Jensen R. Treating tension-type headache: an expert opinion. *Expert Opin Pharmacother* 2011; 12: 1099–109.
10. Andrasik F. Biofeedback in headache: an overview of approaches and evidence. *Cleve Clin J Med* 2010; 77 Suppl 3: S72–6.
11. Bendtsen L, Evers S, Linde M, Mitsikostas DD, Sandrini G, Schoenen J; EFNS.: EFNS guideline on the treatment of tension-type headache-report of an EFNS task force. *Eur J Neurol* 2010; 17: 1318–25.
12. Varkey E, Hagen K, Zwart J-A, Linde M. Physical activity and headache: results from the Nord-Trøndelag Health Study (HUNT). *Cephalalgia* 2008; 28: 1292–7.

14. Wallasch TM, Angeli A, Kropp P. Outcomes of a headache-specific cross-sectional multidisciplinary treatment program. *Headache* 2012; 52: 1094–1105.
12. Varkey E, Hagen K, Zwart J-A, Linde M. Physical activity and headache: results from the Nord-Trøndelag Health Study (HUNT). *Cephalalgia* 2008; 28: 1292–7.
15. Varkey E, Cider A, Carlsson J, Linde M. Exercise as migraine prophylaxis: a randomized study using relaxation and topiramate as controls. *Cephalalgia* 2011; 31: 1428–38.
16. Darabaneanu S, Overath CH, Rubin D, Lühje S, Sye W et al. Aerobic exercise as an option for treating migraine: a pilot study. *Int J Sports Med* 2011; 32: 455–60.
17. Lambert RW Jr, Burnet DL. Prevention of exercise induced-migraine by quantitative warm-up. *Headache* 1985; 25: 317–9.
18. Abbott RB, Hui KK, Hays RD, Li MD, Pan T. A randomized controlled trial of tai chi for tension type headaches. *Evid Based Complement Alternat Med* 2007; 4: 107–13.
19. Köseoglu E, Akboyraz A, Soayuer A, Ersoy AO. Aerobic exercise and plasma beta-endorphin levels in patients with migrainous headache without aura. *Cephalalgia* 2003; 23: 972–6.
20. Varkey E, Cider A, Carlsson J, Linde M. A study to evaluate the feasibility of an aerobic exercise program in patients with migraine. *Headache* 2009; 49: 563–70.
21. Grimm L, Douglas D, Hanson P. Aerobic training in the prophylaxis of migraine. *Med Sci Sports Exerc* 1981; 13: 98.
22. Narin SO, Pinar L, Erbas D, Ozturk V, Idiman F. The effects of exercise and exercise-related changes in blood nitric oxide level on migraine headache. *Clin Rehabil* 2003; 17: 624–30.
23. Gaul C, van Doorn C, Webering N, et al. Clinical outcome of a headache specific multidisciplinary treatment program and adherence to treatment recommendations in a tertiary headache center: An observational study. *J Headache Pain* 2011; 12: 475–483.
24. Gerber W, Miltner W, Gabler H, Hildebrand E, Larbig W. Bewegungs- und Sporttherapie bei chronischen Kopfschmerzen. In: Gerber W, Miltner W, Mayer K (Hrsg.) *Vorergebnisse und Perspektiven interdisziplinärer Forschung*. Weinheim: Edition Medizin 1987: S. 55–66.
25. Nordlander E, Cider A, Carlsson J, Linde M. Improvement of exercise capacity in patients with migraine – Methodological considerations. *Cephalalgia* 2007; 27: 575–759.
26. Fitterling JM, Martin JE, Gramling S, Cole P, Milan MA. Behavioral management of exercise training in vascular headache patients: An investigation of exercise adherence and headache activity. *J Appl Behav Anal* 1988; 21: 9–19.
27. Dittrich SM, Gunther V, Franz G, et al. Aerobic exercise with relaxation: Influence on pain and psychological well-being in female migraine patients. *Clin J Sport Med* 2008; 18: 363–365.
28. Fischer H, Heller R, Giesler A, Hollmann W. Audauertraining bei Migräne: eine Pilotstudie. In: Liesen H, Weiß M, Baum H (Hrsg.) *Regulations- und Repairemechanismen*. 33. Deutscher Sportärztekongress. Köln: Deutscher Ärzteverlag; 1994: S. 807–8.
29. Söderberg EI, Carlsson JY, Stener-Victorin E, Dahlöf C. Subjective well-being in patients with chronic tension-type headache: effect of acupuncture, physical training, and relaxation training. *Clin J Pain* 2011; 27: 448–56.
30. Gunreben-Stempfle B, Griebinger N, Lang E et al. Effectiveness of an intensive multidisciplinary headache treatment program. *Headache* 2009; 49: 900–1000.
31. Domingues RB, Teixeira AL, Domingues SA. Physical practise is associated with less functional disability in medical students with migraine. *Arq Neuropsiquiatr* 2011; 69: 39–43.
32. Sato K, Hayashino Y, Yamazaki S, Takegami S, Ono R et al. Headache prevalence and long working hours: the role of physical inactivity. *Public Health* 2012; 126: 587–93.
33. Dooley JM, Gordon KE, Wood EP, Brna PM. Activity levels among adolescents with Migraine. *Pediatr Neurol* 2006; 35: 119–21.
34. Dias Santos Santiago M, De Souza Carvalho D, Gabbai AA, Machado M, Pinto P, Moutran ARC, Villa TR. Amitriptyline and aerobic exercise or amitriptyline alone in the treatment of chronic migraine: a randomized comparative study. *Arq Neuropsiquiatr* 2014; 72: 851–56.

Literatur zu Kapitel 51 Schlafstörungen

1. Tettenborn B. Schlafstörungen. In: *Prävention und Therapie durch Sport*. Band 2: Neurologie, Psychiatrie/ Psychosomatik, Schmerzsyndrome. 2. Auflage, Elsevier München, 2015.
2. Gerber M, Brand S, Hermann C, Colledge F, Holsboer-Trachsler E, Pühse U. Increased objectively assessed vigorous-intensity exercise is associated with reduced stress, increased mental health and good objective and subjective sleep in young adults. *Physiol Behav* 2014; 135: 17–24.
3. Flausino NH, Prado JM, Queiroz SS, Tufik S, Mello MT. Physical exercise performed before bedtime improves sleep pattern of healthy young good sleepers. *Psychophysiology* 2012; 49: 186–92.

4. Kwon AM, Shin C. Structural equation modelling for the effect of physical exercise on excessive daytime sleepiness. *Public Health* 2016; 141: 95–99.
5. Buman MP, Philips BA, Youngstedt SD, Kline CE, Hirshkowitz M. Does nighttime exercise really disturb sleep? Results from the 2013 National Sleep Foundation Sleep in America Poll. *Sleep Med* 2014; 15: 755–61.
6. Kaczor M, Skalski M. Treatment of behavioural sleep problems in children and adolescents – literature review. *Psychiatr L* 2016; 50: 571–584.
7. Youngdeok K, Masataka U, Lochbaum M, Stegemeier S. Physical activity, screen-based sedentary behaviour, and sleep duration in adolescents: youth risk behaviour survey 2011–2013. *Prev Chronic Dis* 2016; 13: E 131.
8. Buman MP, Hekler EB, Bliwise DL, King AC. Moderators and mediators of exercise induced objective sleep improvements in midlife and older adults with sleep complaints. *Health Psychol* 2011; 30: 579–87.
9. Brand S, Gerber M, Beck J, Hatzinger M, Pühse U, Holsboer-Trachsler E. High exercise levels are related to favorable sleep patterns and psychological functioning in adolescents: a comparison of athletes and controls. *J Adolesc Health* 2010; 46: 133–41.
10. Bonardi JM, Lima LG, Campos GO, Bertani RF, Moriguti JC, Ferriollo E, Lima NKC. Effect of different types of exercise on sleep quality of elderly subjects. *Sleep Medicine* 2016; 25: 122–129.

Literatur zu Kapitel 52 Stress

1. Elfering A, Brunner B, Igic I, Keller A, Weber L. Gesellschaftliche Bedeutung und Kosten von Stress. In: Fuchs R, Gerber M (Hrsg.) *Stressregulation und Sport*. Heidelberg: Springer, 2017.
2. Staatssekretariat für Wirtschaft. *Stressstudie 2010. Stress bei Schweizer Erwerbstätigen. Zusammenhänge zwischen Arbeitsbedingungen, Personenmerkmalen, Befinden und Gesundheit*. Bern: seco, 2011.
3. Semmer NK, Zapf D. Theorien der Stressentstehung und -bewältigung. In: Fuchs R, Gerber M (Hrsg.) *Stressregulation und Sport*. Heidelberg: Springer, 2017.
4. Schwarzer R. *Stress, Angst und Handlungsregulation*. Stuttgart: Kohlhammer, 2000.
5. Gerber M, Fuchs R. *Stressregulation und Sport: Ein Überblick zum Stand der Forschung*. In: Fuchs R, Gerber M (Hrsg.) *Stressregulation und Sport*. Heidelberg: Springer, 2017.
6. Fuchs R, Gerber M (Hrsg.) *Stressregulation und Sport*. Heidelberg: Springer, 2017.
7. Fuchs R, Klaperski S. *Stressregulation durch Sport und Bewegung*. In: Fuchs R, Gerber M (Hrsg.) *Stressregulation und Sport*. Heidelberg: Springer, 2017.
8. Klaperski S. Exercise, stress and health: The stress-buffering effect of exercise. In: Fuchs R, Gerber M (Hrsg.) *Stressregulation und Sport*. Heidelberg: Springer, 2017.
9. Stults-Kolehmainen MA, Sinha R. The effects of stress on physical activity and exercise. *Sports Medicine* 2014; 44: 81–121.
10. Fransson EI, Heikkilä K, Nyberg ST, Zins M, Westerlund H, Westerholm P et al. Job strain as a risk factor for leisure-time physical inactivity: An individual-participant meta-analysis of up to 170'000 men and women. *Am J Epidemiol* 2012; 176: 1078–89.
11. Gerber M, Pühse U. Do exercise and fitness protect against stress-induced health complaints? A review of the literature. *Scandinavian Journal of Public Health* 2009; 37: 801–19.
12. Klaperski S, Seelig H, Fuchs R. Sportaktivität als Stresspuffer. *Zeitschrift für Sportpsychologie* 2012; 19: 80–90.
13. Gerber M, Börjesson M, Ljung T, Lindwall M, Jonsdottir I. Fitness moderates the relationship between stress and cardiovascular risk factors. *Med Sci Sports Exerc* 2016; 48: 2075–81.
14. Gerber M, Endes K, Herrmann C, Colledge F, Brand S, Donath L et al. Fitness, stress and body composition in primary schoolchildren. *Med Sci Sports Exerc* 2017; 49: 581–7.
15. Stults-Kolehmainen MA. The interplay between stress and physical activity in the prevention and treatment of cardiovascular disease. *Front Physiol* 2013; 4: 346.
16. Puterman E, Lin J, Blackburn E, O'Donovan A, Adler N, Epel E. The power of exercise: Buffering the effect of chronic stress on telomere length. *PLoS One* 2010; 5: e10837.
17. Holtermann A, Mortensen OS, Burr H, Søgaard K, Gyntelberg F, Suadicani P. Physical demands at work, physical fitness, and 30-year ischaemic heart disease and all-cause mortality in the Copenhagen Male Study. *Scandinavian Journal of Work and Environmental Health* 2010; 36: 357–65.
18. Sothmann MS. The cross-stressor adaptation hypothesis and exercise training. In: Acevedo EO, Ekkekakis P, editors. *Psychobiology of physical activity*. Champaign: Human Kinetics; 2006. p. 149–60.
19. Gerber M. Physiologische Wirkmechanismen des Sports unter Stress. In: Fuchs R, Gerber M (Hrsg.) *Stressregulation und Sport*. Heidelberg: Springer, 2017.
20. Klaperski S, von Dawans B, Heinrichs M, Fuchs R. Does the level of physical exercise affect physiological and psychological responses to psychosocial stress in women? *Psychology of Sport and Exercise* 2013; 14: 266–74.

21. Klaperski S, von Dawans B, Heinrichs M, Fuchs R. Effects of a 12-week endurance training program on the physiological response to psychosocial stress in men: A randomized controlled trial. *Journal of Behavioral Medicine* 2014; 37: 1118–33.
22. Hamer M, Taylor A, Steptoe A. The effect of acute exercise on stress related blood pressure responses: A systematic review and meta-analysis. *Biological Psychology* 2006; 71: 183–90.
23. Gerber M, Holsboer-Trachsler E, Pühse U, Brand S. Exercise is medicine for patients with major depressive disorders. But only if the “pill” is taken! *Neuropsychiatric Disease and Treatment* 2016; 12: 1977–81.
24. Rosenbaum S, Tiedemann A, Sherrington C, Curtis J, Ward PB. Physical activity interventions for people with mental illness: A systematic review and meta-analysis. *J Clin Psychiatry* 2014; 75: 964–74.